

COMPLEX ANALYSIS¹

Douglas N. Arnold²

References:

John B. Conway, *Functions of One Complex Variable*, Springer-Verlag, 1978.

Lars V. Ahlfors, *Complex Analysis*, McGraw-Hill, 1966.

Raghavan Narasimhan, *Complex Analysis in One Variable*, Birkhäuser, 1985.

CONTENTS

I. THE COMPLEX NUMBER SYSTEM	2
II. ELEMENTARY PROPERTIES AND EXAMPLES OF ANALYTIC FNS.	3
Differentiability and analyticity	4
The Logarithm.....	6
Conformality.....	6
Cauchy–Riemann Equations.....	7
Möbius transformations	9
III. COMPLEX INTEGRATION AND APPLICATIONS TO ANALYTIC FNS.	11
Local results and consequences	12
Homotopy of paths and Cauchy’s Theorem	14
Winding numbers and Cauchy’s Integral Formula.....	15
Zero counting; Open Mapping Theorem	17
Morera’s Theorem and Goursat’s Theorem	18
IV. SINGULARITIES OF ANALYTIC FUNCTIONS.....	19
Laurent series.....	20
Residue integrals.....	23
V. FURTHER RESULTS ON ANALYTIC FUNCTIONS.....	26
The theorems of Weierstrass, Hurwitz, and Montel	26
Schwarz’s Lemma.....	28
The Riemann Mapping Theorem	29
Complements on Conformal Mapping	31
VI. HARMONIC FUNCTIONS	32
The Poisson kernel.....	33
Subharmonic functions and the solution of the Dirichlet Problem	36
The Schwarz Reflection Principle.....	39

¹These lecture notes were prepared for the instructor’s personal use in teaching a half-semester course on complex analysis at the beginning graduate level at Penn State, in Spring 1997. They are certainly not meant to replace a good text on the subject, such as those listed on this page.

²Department of Mathematics, Penn State University, University Park, PA 16802.
Email: dna@math.psu.edu. Web: <http://www.math.psu.edu/dna/>.

I. THE COMPLEX NUMBER SYSTEM

\mathbb{R} is a field. For $n > 1$, \mathbb{R}^n is a vectorspace over \mathbb{R} , so is an additive group, but doesn't have a multiplication on it. We can endow \mathbb{R}^2 with a multiplication by

$$(a, b)(c, d) = (ac - bd, bc + ad).$$

Under this definition \mathbb{R}^2 becomes a field, denoted \mathbb{C} . Note that $(a/(a^2 + b^2), -b/(a^2 + b^2))$ is the multiplicative inverse of (a, b) . (Remark: it is not possible to endow \mathbb{R}^n with a field structure for $n > 2$.) We denote $(0, 1)$ by i and identify $x \in \mathbb{R}$ with $(x, 0)$, so $\mathbb{R} \subset \mathbb{C}$. Thus $(a, b) = a + bi$, $a, b \in \mathbb{R}$. Note that $i^2 = -1$. \mathbb{C} is generated by adjoining i to \mathbb{R} and closing under addition and multiplication. It is remarkable that the addition of i lets us not only solve the equation $x^2 + 1 = 0$, but every polynomial equation.

For a and b real and $z = a + bi$ we define $\operatorname{Re} z = a$, $\operatorname{Im} z = b$, $\bar{z} = a - bi$, and $|z| = (a^2 + b^2)^{1/2}$. Then

$$\begin{aligned} \operatorname{Re} z &= (z + \bar{z})/2, & \operatorname{Im} z &= (z - \bar{z})/(2i), \\ |z|^2 &= z\bar{z}, & \frac{1}{z} &= \frac{\bar{z}}{|z|^2}, \\ \overline{z \pm w} &= \bar{z} \pm \bar{w}, & \overline{z\bar{w}} &= \bar{z}\bar{w}, \\ \overline{z/w} &= \bar{z}/\bar{w}, & |z + w| &\leq |z| + |w|. \end{aligned}$$

The map $\theta \mapsto (\cos \theta, \sin \theta)$ defines a 2π -periodic map of the real line onto the unit circle in \mathbb{R}^2 . In complex notation this map is $\theta \mapsto \operatorname{cis} \theta := \cos \theta + i \sin \theta$. Every nonzero complex number can be written as $r \operatorname{cis} \theta$ where $r > 0$ is uniquely determined and $\theta \in \mathbb{R}$ is uniquely determined modulo 2π . The number 0 is equal to $r \operatorname{cis} \theta$ where $r = 0$ and θ is arbitrary. The relation $z = r \operatorname{cis} \theta$ determines the relations $z \mapsto r$ which is simply the function $r = |z|$ and $z \mapsto \theta$. The latter is denoted $\theta = \arg \theta$. Note that for $z \neq 0$, $\arg \theta$ is determined modulo 2π (while $\arg 0$ is arbitrary). We can normalize \arg by insisting that $\arg z \in (-\pi, \pi]$. Note that if $z_1 = r_1 \operatorname{cis} \theta_1$ and $z_2 = r_2 \operatorname{cis} \theta_2$ then $z_1 z_2 = r_1 r_2 \operatorname{cis}(\theta_1 + \theta_2)$. The latter formula just encapsulates the formula for the sine and cosine of a sum, and gives $\arg z_1 z_2 = \arg z_1 + \arg z_2$. In particular, $ir \operatorname{cis} \theta = r \operatorname{cis}(\theta + \pi/2)$, so multiplication by i is just the operation of rotation by $\pi/2$ in the complex plane. Multiplication by an arbitrary complex number $r \operatorname{cis} \theta$ is just rotation by $\arg \theta$ followed by (or preceded by) dilation by a factor r . Further, $z^n = r^n \operatorname{cis}(n\theta)$. Every nonzero $z \in \mathbb{C}$ admits n distinct n th roots: the n th roots of $r \operatorname{cis} \theta$ are $\sqrt[n]{r} \operatorname{cis}[(\theta + 2\pi k)/n]$, $k = 0, 1, \dots, n$.

Lines and circles in the plane. Circles given by $|z - a| = r$ where $a \in \mathbb{C}$ is the center and $r > 0$ is the radius. If $0 \neq b \in \mathbb{C}$ then the line through the origin in the direction b is the set of all points of the form tb , $t \in \mathbb{R}$, or all z with $\operatorname{Im}(z/b) = 0$. If $t \in \mathbb{R}$ and $c > 0$ then $(t + ci)b = tb + cib$ represents a point in the half plane to the left of b determined by the line tb , i.e., $\{z : \operatorname{Im}(z/b) > 0\}$ is the equation of that half-plane. Similarly, $\{z : \operatorname{Im}[(z - a)/b] > 0\}$ is the translation of that half-plane by a , i.e., the half-plane determined by the line through a parallel to b and in the direction to the left of b .

Stereographic projection determines a one-to-one correspondence between the unit sphere in \mathbb{R}^3 minus the north-pole, S , and the complex plane via the correspondence

$$z \leftrightarrow \frac{x_1 + ix_2}{1 - x_3},$$

$$x_1 = \frac{2 \operatorname{Re} z}{1 + |z|^2}, \quad x_2 = \frac{2 \operatorname{Im} z}{1 + |z|^2}, \quad x_3 = \frac{|z|^2 - 1}{|z| + 1}.$$

If we define $\mathbb{C}_\infty = \mathbb{C} \cup \{\infty\}$, then we have a one-to-one correspondence between S and \mathbb{C}_∞ . This allows us to define a metric on \mathbb{C}_∞ , which is given by

$$d(z_1, z_2) = \frac{2|z_1 - z_2|}{\sqrt{(1 + |z_1|^2)(1 + |z_2|^2)}}, \quad d(z, \infty) = \frac{2}{\sqrt{1 + |z|^2}}.$$

II. ELEMENTARY PROPERTIES AND EXAMPLES OF ANALYTIC FUNCTIONS

For $z \neq 1$, $\sum_{n=0}^N z^n = (1 - z^{N+1})/(1 - z)$. Therefore the geometric series $\sum_{n=0}^{\infty} z^n$ converges (to $1/(1 - z)$) if $|z| < 1$. It clearly diverges, in fact its terms become unbounded, if $|z| > 1$.

Weierstrass M-Test. Let M_0, M_1, \dots be positive numbers with $\sum M_n < \infty$ and suppose that $f_n : X \rightarrow \mathbb{C}$ are functions on some set X satisfying $\sup_{x \in X} |f_n(x)| \leq M_n$. Then $\sum_{n=0}^{\infty} f_n(x)$ is absolutely and uniformly convergent.

Theorem. Let $a_0, a_1, \dots \in \mathbb{C}$ be given and define the number R by

$$\frac{1}{R} = \limsup |a_n|^{1/n}.$$

Then (1) for any $a \in \mathbb{C}$ the power series $\sum_{n=0}^{\infty} a_n(z - a)^n$ converges absolutely for all $|z - a| < R$ and it converges absolutely and uniformly on the disk $|z - a| \leq r$ for all $r < R$. (2) The sequence $a_n(z - a)^n$ is unbounded for all $|z - a| > R$ (and hence the series is certainly divergent).

Thus we see that the set of points where a power series converges consists of a disk $|z - a| < R$ and possibly a subset of its boundary. R is called the radius of convergence of its series. The case $R = \infty$ is allowed.

Proof of theorem. For any $r < R$ we show absolute uniform convergence on $D_r = \{|z - a| \leq r\}$. Choose $\tilde{r} \in (r, R)$. Then, $1/\tilde{r} > \limsup |a_n|^{1/n}$, so $|a_n|^{1/n} < 1/\tilde{r}$ for all n sufficiently large. For such n , $|a_n| < 1/\tilde{r}^n$ and so

$$\sup_{z \in D_r} |a_n(z - a)^n| < (r/\tilde{r})^n.$$

Since $\sum (r/\tilde{r})^n < \infty$ we get the absolute uniform convergence on D_r .

If $|z - a| = r > R$, take $\tilde{r} \in (R, r)$. Then there exist n arbitrarily large such that $|a_n|^{1/n} \geq 1/\tilde{r}$. Then, $|a_n(z - a)^n| \geq (r/\tilde{r})^n$, which can be arbitrarily large. \square

Theorem. If $a_0, a_1, \dots \in \mathbb{C}$ and $\lim |a_n/a_{n+1}|$ exists as a finite number or infinity, then this limit is the radius of convergence R of $\sum a_n(z-a)^n$.

Proof. Without loss of generality we can suppose that $a = 0$. Suppose that $|z| > \lim |a_n/a_{n+1}|$. Then for all n sufficiently large $|a_n| < |a_{n+1}z|$ and $|a_n z^n| < |a_{n+1} z^{n+1}|$. Thus the series $\sum a_n z^n$ has terms of increasing magnitude, and so cannot be convergent. Thus $|z| \geq R$. This shows that $\lim |a_n/a_{n+1}| \geq R$.

Similarly, suppose that $z < \lim |a_n/a_{n+1}|$. Then for all n sufficiently large $|a_n| > |a_{n+1}z|$ and $|a_n z^n| > |a_{n+1} z^{n+1}|$. Thus the series has terms of decreasing magnitude, and so, by the previous theorem, $|z| \leq R$. This shows that $\lim |a_n/a_{n+1}| \leq R$. \square

Remark. On the circle of convergence, many different behaviors are possible. $\sum z^n$ diverges for all $|z| = 1$. $\sum z^n/n$ diverges for $z = 1$, else converges, but not absolutely (this follows from the fact that the partial sums of $\sum z^n$ are bounded for $z \neq 1$ and $1/n \downarrow 0$). $\sum z^n/n^2$ converges absolutely on $|z| \leq 1$. Sierpinski gave a (complicated) example of a function which diverges at every point of the unit circle except $z = 1$.

As an application, we see that the series

$$\sum_{n=0}^{\infty} \frac{z^n}{n!}$$

converges absolutely for all $z \in \mathbb{C}$ and that the convergence is uniform on all bounded sets. The sum is, by definition, $\exp z$.

Now suppose that $\sum_{n=0}^{\infty} a_n(z-a)^n$ has radius of convergence R , and consider its formal derivative $\sum_{n=1}^{\infty} n a_n(z-a)^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1}(z-a)^n$. Now clearly $\sum_{n=0}^{\infty} a_{n+1}(z-a)^n$ has the same radius of convergence as $\sum_{n=0}^{\infty} a_n(z-a)^n$ since

$$(z-a) \sum_{n=0}^N a_{n+1}(z-a)^n = \sum_{n=0}^{N+1} a_n(z-a) - a_0,$$

and so the partial sums on the left and right either both diverge for a given z or both converge. This shows (in a roundabout way) that $\limsup |a_{n+1}|^{1/n} = \limsup |a_n|^{1/n} = 1/R$. Now $\lim(n+1)^{1/n} = 1$ as is easily seen by taking logs. Moreover, it is easy to see that if $\limsup b_n = b$ and $\lim c_n = c > 0$, then $\limsup b_n c_n = bc$. Thus $\limsup |(n+1)a_{n+1}|^{1/n} = 1/R$. This shows that the formal derivative of a power series has the same radius of convergence as the original power series.

Differentiability and analyticity. Definition of differentiability at a point (assumes function is defined in a neighborhood of the point).

Most of the consequences of differentiability are quite different in the real and complex case, but the simplest algebraic rules are the same, with the same proofs. First of all, differentiability at a point implies continuity there. If f and g are both differentiable at a point a , then so are $f \pm g$, $f \cdot g$, and, if $g(a) \neq 0$, f/g , and the usual sum, product, and quotient rules hold. If f is differentiable at a and g is differentiable at $f(a)$, then $g \circ f$ is differentiable at a and the chain rule holds. Suppose that f is continuous at a , g is continuous at $f(a)$, and $g(f(z)) = z$ for all z in a neighborhood of a . Then if $g'(f(a))$ exists and is non-zero, then $f'(a)$ exists and equals $1/g'(f(a))$.

Definition. Let f be a complex-valued function defined on an open set G in \mathbb{C} . Then f is said to be analytic on G if f' exists and is continuous at every point of G .

Remark. We shall prove later that if f is differentiable at every point of an open set in \mathbb{C} it is automatically analytic; in fact, it is automatically infinitely differentiable. This is of course vastly different from the real case.

If Q is an arbitrary non-empty subset of \mathbb{C} we say f is analytic on Q if it is defined and analytic on an open set containing Q .

We now show that a power series is differentiable at every point in its disk of convergence and that its derivative is given by the formal derivative obtained by differentiating term-by-term. Since we know that that power series has the same radius of convergence, it follows that a power series is analytic and infinitely differentiable in its convergence disk. For simplicity, and without loss of generality we consider a power series centered at zero: $f(z) = \sum_n a_n z^n$. Suppose that the radius of convergence is R and that $|z_0| < R$. We must show that for any $\epsilon > 0$, the inequality

$$\left| \frac{f(z) - f(z_0)}{z - z_0} - g(z_0) \right| \leq \epsilon$$

is satisfied for all z sufficiently close to z_0 , where $g(z) = \sum_{n=1}^{\infty} n a_n z^{n-1}$. Let $s_N(z) = \sum_{n=0}^N a_n z^n$, $R_N(z) = \sum_{n=N+1}^{\infty} a_n z^n$. Then

$$\begin{aligned} \left| \frac{f(z) - f(z_0)}{z - z_0} - g(z_0) \right| &\leq \left| \frac{s_N(z) - s_N(z_0)}{z - z_0} - s'_N(z_0) \right| \\ &\quad + |s'_N(z_0) - g(z_0)| + \left| \frac{R_N(z) - R_N(z_0)}{z - z_0} \right| =: T_1 + T_2 + T_3. \end{aligned}$$

Now $s'_N(z_0)$ is just a partial sum for $g(z_0)$, so for N sufficiently large (and all z), $T_2 \leq \epsilon/3$. Also,

$$\frac{R_N(z) - R_N(z_0)}{z - z_0} = \sum_{n=N+1}^{\infty} a_n \frac{z^n - z_0^n}{z - z_0}.$$

Now $|z_0| < r < R$ for some r , and if we restrict to $|z| < r$, we have

$$\left| a_n \frac{z^n - z_0^n}{z - z_0} \right| = |a_n| |z^{n-1} + z^{n-2}z_0 + \cdots + z_0^{n-1}| \leq a_n n r^{n-1}.$$

Since $\sum a_n n r^{n-1}$ is convergent, we have for N sufficiently large and all $|z| < r$ then $T_3 < \epsilon/3$. Now fix a value of N which is sufficiently large by both criteria. Then the differentiability of the polynomial s_N shows that $T_1 \leq \epsilon/3$ for all z sufficiently close to z_0 .

We thus know that if $f(z) = \sum a_n z^n$, then, within the disk of convergence, $f'(z) = \sum n a_n z^{n-1}$, and by induction, $f''(z) = \sum n(n-1) a_n z^{n-2}$, etc. Thus $a_0 = f(0)$, $a_1 = f'(0)$, $a_2 = f''(0)/2$, $a_3 = f'''(0)/3!$, etc. This shows that any convergent power series is the sum of its Taylor series in the disk of convergence:

$$f(z) = \sum \frac{f^n(a)}{n!} (z - a)^n.$$

In particular, $\exp' = \exp$.

Lemma. *A function with vanishing derivative on a region (connected open set) is constant.*

Indeed, it is constant on disks, since we can restrict to segments and use the real result. Then we can use connectedness to see that the set of points where it takes a given value is open and closed.

Use this to show that $\exp(z)\exp(a-z) \equiv \exp a$. Define $\cos z$ and $\sin z$, get $\cos^2 + \sin^2 \equiv 1$, $\exp z = \cos z + i \sin z$, $\operatorname{cis} \theta = \exp(i\theta)$, $\exp z = \exp(\operatorname{Re} z) \operatorname{cis}(\operatorname{Im} z)$.

The Logarithm. If $x \neq 0$, the most general solution of $\exp z = w$ is $z = \log |w| + i \arg w + 2\pi in$, $n \in \mathbb{Z}$. There is no solution to $\exp z = 0$.

Definition. If G is an open set and $f : G \rightarrow \mathbb{C}$ is a continuous function satisfying $\exp(f(z)) = z$, then f is called a branch of the logarithm on G .

Examples: $\mathbb{C} \setminus \mathbb{R}_-$ (principal branch), $\mathbb{C} \setminus \mathbb{R}_+$, a spiral strip.

By the formula for the derivative of an inverse, a branch of the logarithm is analytic with derivative $1/z$.

A branch of the logarithm gives a branch of $z^b := \exp(b \log z)$ (understood to be the principal branch if not otherwise noted). Note that $e^b = \exp b$.

Note that different branches of the logarithm assign values to $\log z$ that differ by addition of $2\pi in$. Consequently different branches of z^b differ by factors of $\exp(2\pi inb)$. If b is an integer, all values agree. If b is a rational number with denominator d there are d values.

Conformality. Consider the angle between two line segments with a common vertex in the complex plane. We wish to consider how this angle is transformed under a complex map. The image will be two smooth curves meeting at a common point, so we have to define the angle between two such curves.

Define a path in the complex plane as a continuous map $\gamma : [a, b] \rightarrow \mathbb{C}$. If $\gamma'(t) \in \mathbb{C}$ exists at some point and is not zero, then it is a vector tangent to the curve at $\gamma(t)$, and hence its argument is the angle between the (tangent to the) curve and the horizontal. (Note: if $\gamma'(t) = 0$, the curve may not have a tangent; e.g., $\gamma(t) = t^2 - it^3$.) Let $\gamma : [a, b] \rightarrow \mathbb{C}$ and $\tilde{\gamma} : [\tilde{a}, \tilde{b}] \rightarrow \mathbb{C}$ be two smooth curves, such that $\gamma(a) = \tilde{\gamma}(\tilde{a}) = z$, and $\gamma'(a), \tilde{\gamma}'(\tilde{a}) \neq 0$. Then $\arg \tilde{\gamma}'(\tilde{a}) - \arg \gamma'(a)$ measures the angle between the curves. Now consider the images under f , e.g., of $\rho(t) = f(\gamma(t))$. Then $\rho'(a) = f'(z)\gamma'(a)$, $\tilde{\rho}'(\tilde{a}) = f'(z)\tilde{\gamma}'(\tilde{a})$. Therefore $\arg \tilde{\rho}'(\tilde{a}) - \arg \rho'(a) = \arg \tilde{\gamma}'(\tilde{a}) + \arg f'(z) - \arg \gamma'(a) - \arg f'(z)$, i.e., the angle is invariant in both magnitude and sign. This says that an analytic mapping is conformal, whenever its derivative is not zero. (Not true if the derivative is zero. E.g., z^2 doubles angles at the origin.)

The conformality of an analytic map arises from the fact that in a neighborhood of a point z_0 such a map behaves to first order like multiplication by $f'(z_0)$. Since $f'(z_0) = r \exp i\theta$, near z_0 f simply behave like rotation by θ followed by dilation. By contrast, a smooth map from $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ can behave like an arbitrary linear operator in a neighborhood of a point.

At this point, show the computer graphics square1.mps and square2.mps.

Cauchy–Riemann Equations. Let $f : \Omega \rightarrow \mathbb{C}$ with $\Omega \subset \mathbb{C}$ open. Abuse notation slightly by writing $f(x, y)$ as an alternative for $f(x + iy)$. If $f'(z)$ exists for some $z = x + iy \in \Omega$, then

$$f'(z) = \lim_{\substack{h \in \mathbb{R} \\ h \rightarrow 0}} \frac{f(z+h) - f(z)}{h} = \lim_{\substack{h \in \mathbb{R} \\ h \rightarrow 0}} \frac{f(x+h, y) - f(x, y)}{h} = \frac{\partial f}{\partial x},$$

and

$$f'(z) = \lim_{\substack{h \in \mathbb{R} \\ h \rightarrow 0}} \frac{f(z+ih) - f(z)}{ih} = \lim_{\substack{h \in \mathbb{R} \\ h \rightarrow 0}} -i \frac{f(x, y+h) - f(x, y)}{h} = -i \frac{\partial f}{\partial y}.$$

Thus complex-differentiability of f at z implies not only that the partial derivatives of f exist there, but also that they satisfy the Cauchy–Riemann equation

$$\frac{\partial f}{\partial x} = -i \frac{\partial f}{\partial y}.$$

If $f = u + iv$, then this equation is equivalent to the system

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}.$$

Another convenient notation is to introduce

$$\frac{\partial f}{\partial z} := \frac{1}{2} \left(\frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} \right), \quad \frac{\partial f}{\partial \bar{z}} := \frac{1}{2} \left(\frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right).$$

(These are motivated by the equations $x = (z + \bar{z})/2$, $y = (z - \bar{z})/(2i)$, which, if z and \bar{z} were independent variables, would give $\partial x/\partial z = 1/2$, $\partial y/\partial z = -i/2$, etc.)

In terms of these, the Cauchy–Riemann equations are exactly equivalent to

$$\frac{\partial f}{\partial \bar{z}} = 0,$$

which is also equivalent to

$$\frac{\partial f}{\partial z} = \frac{\partial f}{\partial x}.$$

Thus, if f is analytic on Ω , then the partial derivatives of f exist and are continuous on Ω , and the Cauchy–Riemann equations are satisfied there. The converse is true as well:

Theorem. *If the partial derivatives of f exist and are continuous on Ω , and the Cauchy–Riemann equations are satisfied there, then f is analytic on Ω .*

Proof. Let $z = x + iy \in \Omega$. We must show that $f'(z)$ exists. Let r be small enough that the disk of radius r around z belongs to Ω and choose $h = s + it$ with $0 < |s + it| < r$. Then, by the mean value theorem,

$$\begin{aligned} \frac{f(z+h) - f(z)}{h} &= \frac{f(x+s, y+t) - f(x, y)}{s+it} \\ &= \frac{f(x+s, y+t) - f(x, y+t)}{s} \frac{s}{s+it} + \frac{f(x, y+t) - f(x, y)}{t} \frac{t}{s+it} \\ &= \frac{\partial f}{\partial x}(x+s^*, y+t) \frac{s}{s+it} + \frac{\partial f}{\partial y}(x, y+t^*) \frac{t}{s+it}, \end{aligned}$$

where $|s^*| < s$ and $|t^*| < t$. Note that

$$\frac{\partial f}{\partial x}(x + s^*, y + t) - \frac{\partial f}{\partial x}(x, y) \rightarrow 0$$

as $h \rightarrow 0$, and similarly for the second partial derivative. Moreover $|s/(s + it)|$ stays bounded (by 1). Thus

$$\frac{f(z + h) - f(z)}{h} = \frac{\partial f}{\partial x}(x, y) \frac{s}{s + it} + \frac{\partial f}{\partial y}(x, y) \frac{t}{s + it} + R(h),$$

where $\lim_{h \rightarrow 0} R(h) = 0$. Now if the Cauchy–Riemann equations hold, then the the first two terms on the right hand side sum to $\partial f / \partial x(x, y)$, which is independent of h , so the limit exists and is equal to $\partial f / \partial x(x, y)$. \square

Remark. The theorem can be weakened to say that if f is continuous on Ω and the partial derivatives exist and satisfy the Cauchy–Riemann equations there (without assuming that the partial derivatives are continuous), then the complex derivative of f exists on Ω (which is equivalent to f being analytic on Ω). This is the Looman–Menchoff Theorem. See Narasimhan, *Complex Analysis in One Variable*, for a proof. We do need at least continuity, since otherwise we could take f to be the characteristic function of the coordinate axes.

Note that

$$\frac{\partial}{\partial z} \frac{\partial}{\partial \bar{z}} = \frac{\partial}{\partial \bar{z}} \frac{\partial}{\partial z} = \frac{1}{4} \Delta.$$

This shows that any analytic function is harmonic (equivalently, its real and imaginary parts are harmonic). It also shows that the conjugate of an analytic function, while not analytic, is harmonic.

If $u(x, y) = \operatorname{Re} f(x + iy)$, then u is harmonic. Conversely, if u is harmonic on a region Ω , does there exist a *harmonic conjugate*, i.e., a function v on Ω such that $f = u + iv$ is analytic? Clearly, if there exists v it is determined up to addition of a real constant (or f is determined up to addition of an imaginary constant). The book gives an elementary construction of the harmonic conjugate in a disk and in the whole plane. We sketch the idea of a more general proof. We require that the domain satisfy the condition that if γ is any piecewise simple smooth closed curve in Ω , then γ is the boundary of a subset G of Ω . In other words, Ω has no holes (is simply-connected). Our proof follows directly from the following result, which holds on simply-connected domains: let $f : \Omega \rightarrow \mathbb{R}^2$ be a C^1 vectorfield. Then there exists $v : \Omega \rightarrow \mathbb{R}$ such that $f = \nabla v$ if and only if $\partial f_1 / \partial y = \partial f_2 / \partial x$. The “only if” part is obvious. To prove the “if” part, fix a point (x_0, y_0) in Ω and for any $(x, y) \in \Omega$, let $\gamma_{(x,y)}$ be a piecewise smooth path in Ω from (x_0, y_0) to (x, y) , and let τ be the unit tangent to the path pointing from (x_0, y_0) towards (x, y) . Define

$$v(x, y) = \int_{\gamma_{(x,y)}} f \cdot \tau \, ds.$$

It is essential that this quantity doesn't depend on the choice of path, or, equivalently, that

$$\int_{\gamma} f \cdot \tau ds = 0$$

for all piecewise smooth simple closed paths γ . But

$$\int_{\gamma} f \cdot \tau ds = \int_{\gamma} (f_2, -f_1) \cdot n ds = \iint \nabla \cdot (f_2, -f_1) dx dy = 0,$$

by the divergence theorem. It is easy to check that $\nabla v = f$.

For a non-simply connected region, there may exist no harmonic conjugate. E.g., if $u = \log |z|$ on $\mathbb{C} \setminus \{0\}$, then $u + i \arg z$ is analytic on $\mathbb{C} \setminus \{z \leq 0\}$, but cannot be extended to an analytic function on $\mathbb{C} \setminus \{0\}$.

Möbius transformations.

Definition. Let a, b, c, d be complex numbers with $ad \neq bc$. Then the mapping

$$S(z) = \frac{az + b}{cz + d}, \quad z \in \mathbb{C}, \quad z \neq -d/c,$$

is a Möbius transformation.

Remarks. 1) Note that if $ad = bc$ the same expression would yield a constant. 2) The coefficients aren't unique, since we can multiply them all by any nonzero complex constant. To each MT we associate the nonsingular matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ of its coefficients, which is determined up to a non-zero multiple. 3) The linear (but non-constant) polynomials are MTs, namely the ones for which $c = 0$. 4) If $z \neq -d/c$ then $S(z) \in \mathbb{C}$ and $S'(z) = (ad - bc)/(cz + d)^2 \neq 0$. 5) We can define $S(-d/c) = \infty$ and $S(\infty) = a/c$, so S can be viewed of as a map from \mathbb{C}_{∞} into itself (which is, as we will now see, 1-1 and onto).

If S and T are MTs, then so is $S \circ T$, its coefficient matrix being the product of the coefficient matrices of S and T . Any MTs admits an inverse, namely the transform with the inverse coefficient matrix. With the understanding that $S(\infty) = a/c$ and $S(-d/c) = \infty$ (or, if $c = 0$, then $S(\infty) = \infty$, the MT maps \mathbb{C}_{∞} 1-1 onto itself. This is evident from the existence of its inverse $S^{-1}(z) = (dz - b)/(-cz + a)$. Moreover, the composition of Möbius transformations is again one, so they form a group under composition. If S is a non-linear MT we can write it as

$$\frac{az + b}{z + d} = a + \frac{b - ad}{z + d}.$$

This shows that any MT can be written as a composition of translations, rotations and dilations (multiplication by a complex number), and inversion (reciprocals).

Since the geometric action of translation, dilation, and rotation are quite evident, let's consider inversion. It takes the unit disk D to $\mathbb{C}_{\infty} \setminus \bar{D}$. We now show that inversion maps lines and circles to other lines and circles.

Consider the equation $|z - p| = k|z - q|$ where $k > 0$, $p, q \in \mathbb{C}$. This is

$$(x - a)^2 + (y - b)^2 = k^2[(x - c)^2 + (y - d)^2]$$

which is clearly the equation of a circle if $k \neq 1$ and the equation of a line if $k = 1$. In either case we have a circle in the Riemann sphere, which we call C_1 . Substituting $1/z$ for z and doing some simple manipulations gives the equation $|z - 1/p| = (k|q|/|p|)|z - 1/q|$, which is another such circle, C_2 . Thus we have $1/z \in C_1 \iff z \in C_2$, showing that inversion maps circles in the sphere onto other circles in the sphere. Since the same property is evident for translation, dilation, and rotation, we conclude that this property holds for all MT.

Some further analysis, which will be omitted, establishes two further properties. 1) A MT is orientation preserving in the sense that, if we traverse a circle in the order of three distinct points on it, z_1, z_2, z_3 , the region to the left of the circle will map to the region to the left of the image circle, with respect to the image orientation. 2) A MT preserves the property of two points being symmetric with respect to a circle, i.e., lying on the same ray from the center, and such that the geometric mean of their distances from the center equals the radius.

Let z_2, z_3, z_4 be distinct points in \mathbb{C} . Then it is easy to see that there is a unique MT taking these points to 1, 0, and ∞ , respectively, namely

$$Sz = \frac{z - z_3}{z - z_4} \frac{z_2 - z_4}{z_2 - z_3}.$$

We can also handle, as a special case, the possibility that one of the z_i is ∞ . We infer from this and the invertibility of the MTs that given two ordered triples of distinct points in \mathbb{C}_∞ , there is a unique MT taking the first onto the second.

If we have any two disks or halfplanes, or one of each, we can map one to the other by a MT, and can arrange that the image of any three points on the boundary of one go to three points on the boundary of the other.

The book makes heavy use of the notation (z_1, z_2, z_3, z_4) (cross ratio) for the image of z_1 under the transformation taking z_2, z_3, z_4 to 1, 0, and ∞ , respectively, so

$$(z_1, z_2, z_3, z_4) = \frac{z_1 - z_3}{z_1 - z_4} \frac{z_2 - z_4}{z_2 - z_3}.$$

From the definition, $(Tz_1, Tz_2, Tz_3, Tz_4) = (z_1, z_2, z_3, z_4)$ for any MT T .

Example: $z \mapsto (1 - z)/(1 + z)$ takes the right half plane onto the unit disk.

III. COMPLEX INTEGRATION AND APPLICATIONS TO ANALYTIC FUNCTIONS

If $\gamma : [a, b] \rightarrow \mathbb{C}$ is piecewise smooth, and f is a continuous complex-valued function defined on the image of γ , define

$$\int_\gamma f(z) dz = \int_a^b f(\gamma(t))\gamma'(t) dt.$$

This is simply the line integral of f along γ . It can be defined analogously to the Riemann integral as the limit of sums of the form $\sum (f \circ \gamma)(\tau_k)[\gamma(t_k) - \gamma(t_{k-1})]$, so is the *Riemann–Stieltjes integral* of $f \circ \gamma$ with respect to γ . Using this definition it can be extended to rectifiable paths, i.e., ones for which γ is only of bounded variation.

If ϕ is a piecewise smooth increasing function from $[c, d]$ onto $[a, b]$ and we let $\rho = \gamma \circ \phi$ (so ρ is the same path as γ but with a different parametrization), then $\int_\rho f(z) dz = \int_\gamma f(z) dz$.

From the fundamental theorem of calculus we see that if F is any analytic function on a neighborhood of γ , then $\int_\gamma F'(z) dz = F(\beta) - F(\alpha)$ where $\alpha = f(a)$, $\beta = f(b)$. If γ is a closed path, this is zero.

As a consequence, if $\gamma = re^{i\theta}$, then

$$\int_\gamma z^n dz = 0 \quad \text{for all } n \in \mathbb{Z}, n \neq -1.$$

By direct calculation we have $\int_\gamma z^{-1} dz = 2\pi i$.

More generally, let $|z_0| < r$ and consider $\int_\gamma \frac{1}{z-z_0} dz$. Let z_ϵ^\pm be the point on the circle $|z - z_0| = r$ with $\arg(z_\epsilon^\pm - z_0) = \pm(\pi - \epsilon)$. Clearly, as $\epsilon \rightarrow 0$, $z_\epsilon^\pm \rightarrow z_1$, the point on the circle with the same imaginary part as z_0 and negative real part. Now let γ_ϵ be the curve running from z_ϵ^- to z_ϵ^+ on the circle. In a neighborhood of γ_ϵ the function $\text{Log}(z - z_0)$ is a primitive of $1/(z - z_0)$. So

$$\int_{\gamma_\epsilon} \frac{1}{z - z_0} dz = \text{Log}(z_\epsilon^+ - z_0) - \text{Log}(z_\epsilon^- - z_0) = \log r + i(\pi - \epsilon) - \log r + i(\pi - \epsilon).$$

Taking the limit as $\epsilon \downarrow 0$ gives

$$\int_\gamma \frac{1}{z - z_0} dz = 2\pi i.$$

Local results and consequences.

Theorem (Cauchy's integral formula for a disk). *Let f be analytic on a closed disk and let γ be a path around the boundary of the disk. Then*

$$f(z) = \frac{1}{2\pi i} \int_\gamma \frac{f(w)}{w - z} dw$$

for all z in the open disk.

Proof. Consider the function

$$g(t) = \int_\gamma \frac{f(z + t(w - z))}{w - z} dw.$$

Then

$$g'(t) = \int_{\gamma} f'(z + t(w - z)) dw,$$

and, since $F(w) = f(z + t(w - z))/t$ is a primitive for $f'(z + t(w - z))$, $g'(t) \equiv 0$. Therefore,

$$\int_{\gamma} \frac{f(w)}{w - z} dw = g(1) = g(0) = f(z) \int_{\gamma} \frac{1}{w - z} dw = 2\pi i f(z). \quad \square$$

As an easy corollary, we have:

Cauchy's Theorem for a disk. *Let f be analytic on a closed disk and let γ be a path around the boundary of the disk. Then*

$$\int_{\gamma} f(w) dw = 0.$$

Proof. Fix z in the open disk and apply Cauchy's formula to the function $w \mapsto f(w)(w - z)$. \square

We can now show that an analytic (i.e., continuously complex differentiable) function has a power series expansion around each point in its domain. (And we already know the converse is true.)

Theorem. *Let f be analytic in $B(a; R)$. Then $f(z) = \sum a_n(z - a)^n$ for $|z - a| < R$, for some $a_n \in \mathbb{C}$.*

Proof. Fix z and choose $r < R$ with $|z - a| < r$. Let γ be the path around $|w - a| = r$. Then $|(z - a)/(w - a)| < 1$, so the Weierstrass M-test implies that the series

$$\sum \frac{1}{(w - a)^{n+1}} (z - a)^n = \frac{1}{w - z},$$

converges absolutely and uniformly in w for $|w - a| = r$. Since f is bounded on the circle, the same sort of convergence holds for

$$\sum \frac{f(w)}{(w - a)^{n+1}} (z - a)^n = \frac{f(w)}{w - z}.$$

Therefore we can interchange summation and integration to get

$$f(z) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(w)}{w - z} dw = \sum a_n (z - a)^n,$$

where

$$a_n = \int_{\gamma} \frac{f(w)}{(w - a)^{n+1}} dw. \quad \square$$

Corollary (of proof). *Under the same hypotheses*

$$f^{(n)}(a) = \frac{n!}{2\pi i} \int_{\gamma} \frac{f(w)}{(w-a)^{n+1}} dw.$$

Corollary (Cauchy's estimate). *If f is analytic in $B(a; R)$, then*

$$|f^{(n)}(a)| \leq \frac{n! \max_{B(a; R)} |f|}{R^n}.$$

Proof. Evident if R is replaced by $r < R$. Then pass to limit. \square

Corollary (Maximum Modulus Principle). *An analytic function on a region which achieves a maximum modulus in the region is constant.*

Proof. By the Cauchy estimate for $n = 0$, if the function achieves its maximum modulus at a point, it achieves it everywhere on a disk containing the point. Thus, the set of points where the maximum modulus is achieved is open. It is also clearly closed. Hence the function has constant modulus. It follows from the Cauchy–Riemann equations that the function itself is constant. \square

Corollary (Liouville's Theorem). *A bounded entire function is constant.*

Proof. Cauchy's estimate for $n = 1$ implies $f' \equiv 0$. \square

Corollary (Fundamental Theorem of Algebra). *Every nonconstant polynomial has a root in \mathbb{C} .*

Theorem. *If f is analytic on a region then there does not exist a point a such that $f^{(n)}(a) = 0$ for all n , unless f is identically zero on the region.*

Proof. The set of such a is closed, and using the Taylor expansion, is open. \square

Corollary. *If f is analytic on a region Ω and not identically zero, then for each $a \in \Omega$ there exists a unique integer $p \in \mathbb{N}$ and a function g , analytic in a neighborhood of a and nonzero at a , such that*

$$f(z) = (z-a)^p g(z).$$

Proof. $f(z) = \sum c_n (z-a)^n$ with $c_n = f^{(n)}(a)/n!$. Let p be the least integer with $c_p \neq 0$ (which exists by the preceding theorem), and $g(z) = \sum_{n=p}^{\infty} c_n (z-a)^{n-p}$. \square

The number p is the *multiplicity* of a as a root of f .

Corollary. *If f is analytic on a region and not identically zero, then the roots of f are isolated.*

Corollary. *An analytic function on a region is completely determined by its value on any set of points containing a limit point.*

Homotopy of paths and Cauchy's Theorem.

Definition. Let $\gamma_0, \gamma_1 : [0, 1] \rightarrow G$ be two closed paths in a region $G \subset \mathbb{C}$. We say that the paths are *homotopic in G* if there exists a continuous function $\Gamma : [0, 1] \times [0, 1] \rightarrow G$ such that for each s $\Gamma(\cdot, s)$ is a closed path with $\Gamma(\cdot, 0) = \gamma_0$ and $\Gamma(\cdot, 1) = \gamma_1$.

This is an equivalence relation, written $\gamma_0 \sim \gamma_1$ in G .

Theorem. Suppose that γ_0 and γ_1 are two homotopic piecewise smooth closed curves in G and that f is analytic on G . Then

$$\int_{\gamma_0} f(z) dz = \int_{\gamma_1} f(z) dz.$$

If we assume that the homotopy Γ is C^2 , the result is easy: let

$$g(s) = \int_0^1 f(\Gamma(t, s)) \frac{\partial \Gamma}{\partial t}(t, s) dt.$$

Then

$$g'(s) = \int_0^1 \frac{\partial}{\partial s} [f(\Gamma(t, s)) \frac{\partial \Gamma}{\partial t}(t, s)] dt = \int_0^1 \frac{\partial}{\partial t} [f(\Gamma(t, s)) \frac{\partial \Gamma}{\partial s}(t, s)] dt = 0,$$

since both integrands equal

$$f'(\Gamma(s, t)) \frac{\partial \Gamma}{\partial t} \frac{\partial \Gamma}{\partial s} + f(\Gamma(s, t)) \frac{\partial^2 \Gamma}{\partial t \partial s}.$$

The proof without assuming smoothness is based on Cauchy's theorem for a disk and a polygonal approximation argument.

Sketch of proof. Since Γ is continuous on the compact set $[0, 1] \times [0, 1]$, it is uniformly continuous. Moreover, there exists $\epsilon > 0$ so that G contains a closed disk of radius ϵ about each point in the range of Γ . Therefore, we can find an integer n , so that Γ maps each of the squares $[j/n, (j+1)/n] \times [k/n, (k+1)/n]$ into the disk of radius ϵ about $\Gamma(j/n, k/n)$. Let ρ_k be the closed polygonal path determined by the points $\Gamma(j/n, k/n)$, $j = 0, 1, \dots, n$. It is easy to conclude from Cauchy's theorem for a disk that

$$\int_{\rho_k} f(z) dz = \int_{\rho_{k+1}} f(z) dz,$$

and also that

$$\int_{\gamma_0} f(z) dz = \int_{\rho_0} f(z) dz, \quad \int_{\gamma_1} f(z) dz = \int_{\rho_n} f(z) dz. \quad \square$$

Definition. A closed path in G is *homotopic to 0* in G if it is homotopic to a constant path.

Corollary. If γ is a piecewise smooth path which is homotopic to 0 in G , and f is analytic in G , then

$$\int_{\gamma} f(z) dz = 0.$$

Definition. A region G is simply-connected if every closed path in G is homotopic to 0.

As an immediate consequence of the theorem above, we have Cauchy's theorem (previously proved only for G a disk).

Cauchy's Theorem. If G is a simply-connected region in the complex plane then

$$\int_{\gamma} f(z) dz = 0$$

for every function f analytic in G and every piecewise smooth closed curve γ in G .

Winding numbers and Cauchy's Integral Formula. Intuitive meaning of winding number: if a pole is planted in the plane at a point a and a closed curve is drawn in the plane not meeting a , if the curve were made of well-lubricated rubber and were to be contracted as small as possible, if it contracts to a point other than a its winding number is 0. If it contracts around the pole planted at a , its winding number is the number of times it circles the pole (signed by orientation).

Definition. Let γ be a piecewise smooth closed curve in \mathbb{C} and $a \in \mathbb{C} \setminus \gamma$. Then the *index* of γ with respect to a , or the *winding number* of γ about a is defined to be

$$n(\gamma; a) = \frac{1}{2\pi i} \int_{\gamma} \frac{1}{z - a} dz.$$

Demonstrate how the winding number can be calculated by drawing a negatively directed ray from a and integrating over pieces of the curve between successive intersections of the curve with the ray.

Proposition. The index $n(\gamma; a)$ is an integer for all $a \in \mathbb{C} \setminus \gamma$. It is constant on each connected component of $\mathbb{C} \setminus \gamma$.

Proof. Assuming γ is parametrized by $[0, 1]$, define

$$g(t) = \int_0^t \frac{\gamma'(s)}{\gamma(s) - a} ds.$$

An easy direct calculation shows that

$$\frac{d}{dt} \exp[-g(t)][\gamma(t) - a] = 0,$$

so, indeed, $\exp[-g(t)][\gamma(t) - a] = \gamma(0) - a$. Taking $t = 1$,

$$\exp[-2\pi i n(\gamma; a)][\gamma(1) - a] = \gamma(0) - a,$$

or, $\exp[-2\pi i n(\gamma; a)] = 1$, which implies that $n(\gamma; a)$ is indeed integral.

The function is obviously continuous, and being integer, it is constant on connected components. Clearly also, it tends to 0 as a tends to ∞ , so it is identically 0 on the unbounded component. \square

The next result is an immediate corollary of the invariance of path integrals of analytic functions under homotopy.

Proposition. *If γ_0 and γ_1 are paths which are homotopic in $\mathbb{C} \setminus \{a\}$ for some point a , then $n(\gamma_0; a) = n(\gamma_1; a)$.*

Cauchy's Integral Formula. *Let γ a piecewise smooth curve in a region G which is null homotopic there, and let f be an analytic function on G . Then*

$$n(\gamma; a)f(a) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{z - a} dz, \quad a \in G \setminus \gamma.$$

Proof. For $a \in G \setminus \gamma$ fixed, let

$$g(z) = \begin{cases} \frac{f(z) - f(a)}{z - a}, & z \in G \setminus \{a\}, \\ f'(a), & z = a. \end{cases}$$

It is easy to show that g is continuous (even analytic!) on G (show the continuity as a homework exercise). Applying Cauchy's theorem to g gives the desired result immediately.

Remark. This presentation is quite a bit different from the book's. The book proves Cauchy's formula under the assumption that γ is *homologous to 0*, i.e., that $n(\gamma; a) = 0$ for all $a \in \mathbb{C} \setminus G$. It is easy to see that homotopic to 0 implies homologous to 0, but the reverse is not true, so the book's result is a bit stronger. It is true that all piecewise smooth closed curves in region are homologous to zero iff G is simply-connected (see Theorem 2.2 of Chapter 8).

The book also goes on to consider chains, which are unions of curves and proves that

$$f(a) \sum_{j=1}^k n(\gamma_j; a) = \frac{1}{2\pi i} \sum_{j=1}^k \int_{\gamma_j} \frac{f(w)}{w - z} dz,$$

if the chain $\gamma_1 + \cdots + \gamma_n$ is homologous to zero, i.e., $\sum_{j=1}^k n(\gamma_j; z) = 0$ for all $z \in \mathbb{C} \setminus G$. This sort of result is useful, e.g., in the case of an annulus. However we can deal with this solution in other ways (by adding a line connecting the circles).

Zero counting; Open Mapping Theorem.

Theorem. Let G be a region, γ a curve homotopic to 0 in G and f an analytic function on G with zeros a_1, a_2, \dots, a_m repeated according to multiplicity. Then

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = \sum_{k=1}^m n(\gamma; a_k).$$

Proof. We can write $f(z) = (z - a_1)(z - a_2) \dots (z - a_m)g(z)$ where $g(z)$ is analytic and nonzero on G . Then

$$\frac{f'(z)}{f(z)} = \frac{1}{z - a_1} + \frac{1}{z - a_2} + \dots + \frac{1}{z - a_m} + \frac{g'(z)}{g(z)},$$

and the theorem follows. \square

If the a_i are the solutions of the equation $f(z) = \alpha$, then

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z) - \alpha} dz = \sum_{k=1}^m n(\gamma; a_k).$$

Note that we may as well write the integral as

$$\frac{1}{2\pi i} \int_{f \circ \gamma} \frac{1}{w - \alpha} dw.$$

Theorem. If f is analytic at a and $f(a) = \alpha$ with finite multiplicity $m \geq 1$, then there exists $\epsilon, \delta > 0$ such that for all $0 < |\zeta - \alpha| < \delta$, the equation $f(z) = \zeta$ has precisely m roots in $|z - a| < \epsilon$ and all are simple.

Proof. That the number of roots is m comes from the continuity of the integral as long as ζ remains in the same connected component of $\mathbb{C} \setminus f \circ \gamma$ as α , and the fact that it is integer valued. We can insure that the roots are simple by taking ϵ small enough to avoid a root of f' . \square

Open Mapping Theorem. A non-constant analytic function is open.

Proof. This is an immediate corollary.

Morera's Theorem and Goursat's Theorem.

Morera's Theorem. *If G is a region and $f : G \rightarrow \mathbb{C}$ a continuous function such that $\int_T f dz = 0$ for each triangular path T in G , then f is analytic.*

Proof. Since it is enough to show f is analytic on disks, we can assume G is a disk. Let a be the center of the disk, and define $F(z) = \int_{[a,z]} f$. By hypothesis

$$\frac{F(z) - F(z_0)}{z - z_0} = \frac{1}{z - z_0} \int_{[z_0,z]} f(w) dw = \int_0^1 f(z_0 + t(z - z_0)) dt.$$

Continuity then implies that $F'(z_0) = f(z_0)$, so f has a primitive, so is analytic.

Before proceeding to Goursat's Theorem, we consider another result that uses an argument similar to Morera's Theorem.

Theorem. *Let G be a region and $f : G \rightarrow \mathbb{C}$ a continuous function. Then f admits a primitive on G if and only if $\int_\gamma f = 0$ for all piecewise smooth closed curves γ in G .*

Proof. Only if is clear. For the if direction, fix $a \in G$ and define $F(z) = \int_{\gamma(a,z)} f$ where $\gamma(a, z)$ is any piecewise smooth path from a to z . Reasoning as in Morera's theorem $F' = f$. \square

In particular, if G is a simply-connected region then every analytic function admits a primitive. If $0 \notin G$, then $1/z$ admits a primitive. Adjusting the primitive by a constant we can assure that it agrees with a logarithm of z_0 at some point z_0 . It is then easy to check by differentiation that $\exp(F(z))/z \equiv 1$. Thus

Proposition. *If G is a simply connected domain, $0 \in G$, then a branch of the logarithm may be defined there.*

(As an interesting example, take G to be the complement of the spiral $r = \theta$.)

We can at long last show that a function which is complex differentiable in a region (but without assuming it is continuously differentiable) is analytic.

Goursat's Theorem. *Let G be a region and $f : G \rightarrow \mathbb{C}$ complex differentiable at each point of G . Then f is analytic.*

Proof. We will show that the integral of f vanishes over every triangular path. Let Δ_0 be a closed triangle with boundary T_0 . Using repeated quadrisection we obtain triangles

$$\Delta_0 \subset \Delta_1 \subset \dots$$

with boundaries T_n such that

$$\left| \int_{T_0} f \right| \leq 4^n \left| \int_{T_n} f \right|.$$

Let $d = \text{diam } T_0$, $p = \text{perim } T_0$. Then $\text{diam } T_n = d/2^n$, $\text{perim } T_n = p/2^n$. By compactness $\bigcap \Delta_n \neq \emptyset$. Let z_0 be in the intersection (it's unique).

From the differentiability of f at z_0 we have: given $\epsilon > 0$

$$|f(z) - f(z_0) - f'(z_0)(z - z_0)| \leq \epsilon |z - z_0|$$

for all z in some disk around z_0 , therefore for all $z \in \Delta_n$, some n . Since $\int_{T_n} f(z) dz = \int_{T_n} z dz = 0$,

$$\begin{aligned} \left| \int_{T_n} f \right| &= \left| \int_{T_n} [f(z) - f(z_0) - f'(z_0)(z - z_0)] dz \right| \\ &\leq \epsilon \text{diam } T_n \text{perim } T_n = \epsilon dp/4^n. \end{aligned}$$

Therefore $\left| \int_{T_0} f \right| \leq \epsilon dp$, and, since ϵ was arbitrary, it must vanish. \square

IV. SINGULARITIES OF ANALYTIC FUNCTIONS

Definition. If f is analytic in $B(a; R) \setminus \{a\}$, for some $a \in \mathbb{C}$, $R > 0$, but not in $B(a; R)$. We say f has an *isolated singularity* at a . If there exists g analytic on some disk centered at a which agrees with f except at a , we say f has a *removable singularity*.

Theorem. Suppose f has an isolated singularity at a . It is removable if and only if $\lim_{z \rightarrow a} (z - a)f(z) = 0$.

Proof. “Only if” is clear. For “if” let $g(z) = (z - a)f(z)$, $z \neq a$, $g(a) = 0$. Using Morera’s theorem (separating the cases a inside the triangle, on the boundary, or in the interior), it is easy to see that g is analytic. Therefore $g(z) = (z - a)h(z)$ for h analytic, and h is the desired extension of f . \square

Corollary. If $\lim_{z \rightarrow a} f(z) \in \mathbb{C}$, or even $f(z)$ is bounded in a punctured neighborhood of a , or even $|f(z)| \leq C|z - a|^{-(1-\epsilon)}$ for some $\epsilon > 0$, some C , then f has a removable singularity.

Now either $\lim_{z \rightarrow a} |f(z)|$ exists as a finite real number, equals $+\infty$, or does not exist as a finite number or infinity. The first case is that of a removable singularity. The second is called a *pole*, the third, an *essential singularity*.

Note that if f has a pole at a , then $1/f(z)$ has a removable singularity there and extends to a with value 0. Using this we see that f has a pole at a if and only if $f(z) = (z - a)^{-m}g(z)$ for some integer $m > 0$ and g analytic in a neighborhood of a . Thus

$$f(z) = \sum_{n=-m}^{\infty} a_n (z - a)^n$$

in a punctured neighborhood of a .

The function $e^{1/z}$ has an essential singularity since

$$\lim_{z \downarrow 0} e^{1/z} = +\infty, \quad \lim_{z \uparrow 0} e^{1/z} = 0.$$

Theorem (Casorati-Weierstrass). *If f has an essential singularity at a , then the image under f of any punctured disk around a is dense in \mathbb{C} .*

Proof. WLOG, assume $a = 0$. If the conclusion is false, there is a punctured disk around 0 in which $f(z)$ stays a fixed positive distance ϵ away from some number $c \in \mathbb{C}$. Consider the function $g(z) = [f(z) - c]/z$. It tends to ∞ as $z \rightarrow 0$, so it has a pole at 0. Therefore $z^m[f(z) - c] \rightarrow 0$ as $z \rightarrow 0$ for sufficiently large m . Therefore, $z^m f(z) \rightarrow 0$, which is not possible if f has an essential singularity.

Laurent series.

Definition. Suppose f is analytic on $\{z : |z| > R\}$ for some R . Then we say that f has an isolated singularity at ∞ . It is removable if $f(1/z)$ has a removable singularity at 0.

We easily see that f has a removable singularity at $\infty \iff \exists \lim_{z \rightarrow \infty} f(z) \in \mathbb{C} \iff f$ is bounded in a neighborhood of infinity.

Note that if f has a removable singularity at ∞ then we can expand f in a power series in $1/z$, convergent for $|z| > R$ uniformly and absolutely for $|z| \geq \rho > R$.

We say that a series of the form $\sum_{n=-\infty}^{\infty} a_n(z-a)^n$ converges if both the series $\sum_{n=1}^{\infty} a_{-n}(z-a)^{-n}$ and $\sum_{n=0}^{\infty} a_n(z-a)^n$ converge (and their sum is the value of the doubly infinite summation).

Lemma. *Let γ be a piecewise smooth curve and g a continuous complex-valued function on γ . Define*

$$f(z) = \int_{\gamma} \frac{g(w)}{z-w} dw, \quad z \notin \gamma.$$

The f is analytic on $\mathbb{C} \setminus \gamma$.

Proof. Fix $z \notin \gamma$. Then

$$\frac{\frac{g(w)}{\zeta-w} - \frac{g(w)}{z-w}}{\zeta-z} = -\frac{g(w)}{(\zeta-w)(z-w)} \rightarrow -\frac{g(w)}{(z-w)^2}$$

as $\zeta \rightarrow z$ with the convergence uniform in $w \in \gamma$. Thus f is differentiable at each $z \in \mathbb{C} \setminus \gamma$. \square

Let f be defined and analytic in an annulus $R_1 < |z| < R_2$. Define

$$f_2(z) = \frac{1}{2\pi i} \int_{|w|=r} \frac{f(w)}{w-z} dw,$$

where $R_2 > r > |z|$ (it doesn't matter which such r we take, by Cauchy's theorem). Then f_2 is analytic in $|z| < R_2$ and so has a convergent power series there:

$$f_2(z) = \sum_{n=0}^{\infty} a_n z^n.$$

Also, let

$$f_1(z) = -\frac{1}{2\pi i} \int_{|w|=r} \frac{f(w)}{w-z} dw$$

where now r is taken with $R_1 < r < |z|$. Then this is analytic in $|z| > R_1$ and tends to 0 at ∞ . Thus

$$f_1(z) = \sum_{n=1}^{\infty} a_{-n} z^{-n}.$$

Now, it is easy to use Cauchy's integral formula to see that $f = f_1 + f_2$ in the annulus. This shows that

$$f(z) = \sum_{n=-\infty}^{\infty} a_n z^n,$$

with the convergence absolute and uniform on compact subsets of the annulus. Note that such an expansion is unique, since we can easily recover the a_n :

$$a_m = \frac{1}{2\pi i} \sum_{n \in \mathbb{Z}} a_n \int_{|z|=r} \frac{z^n}{z^{m+1}} dz = \frac{1}{2\pi i} \int_{|z|=r} \frac{f(z)}{z^{m+1}} dz$$

where r is any number in (R_1, R_2) (result is independent of choice of r since two different circles are homotopic in the annulus).

Summarizing:

Theorem (Laurent Expansion). *If f is an analytic function on the annulus $G = \text{ann}(0; R_1, R_2)$ for some $0 \leq R_1 < R_2 \leq \infty$, then*

$$f(z) = \sum_{n=-\infty}^{\infty} a_n z^n$$

for all z in the annulus. The convergence is uniform and absolute on compact subsets of the annulus and the coefficients a_n are uniquely determined by the formula above.

If f has an isolated singularity at a , it has a Laurent expansion in a punctured disk at a :

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z-a)^n, \quad 0 < |z-a| < R.$$

We define

$$\text{ord}_a(f) = \inf\{n \in \mathbb{Z} : a_n \neq 0\}.$$

Thus f has a removable singularity $\iff \text{ord}_a(f) \geq 0$, and it extends to a as 0 $\iff \text{ord}_a(f) > 0$, in which case this is the order of the 0. $\text{ord}_a(f) = +\infty \iff f$ is identically 0 near a . $\text{ord}_a(f) = -\infty \iff f$ has an essential singularity at a . $\text{ord}_a(f) < 0$ but finite $\iff f$ has a pole, in which case $-\text{ord}_a(f)$ is the order of the pole.

We call a_{-1} the *residue* of f at a , denoted $\text{Res}(f; a)$. Note that, if f has an isolated singularity at a , then for r sufficiently small,

$$\text{Res}(f; a) = \frac{1}{2\pi i} \int_{|z-a|=r} f(z) dz.$$

Give pictorial proof that if G is simply-connected domain, f analytic on G except for isolated singularities, and γ is a simple closed curve that encloses singularities at a_1, a_2, \dots, a_m (necessarily finite), then

$$\frac{1}{2\pi i} \int_{\gamma} f(z) dz = \sum_{j=1}^m \text{Res}(f; a_j).$$

a simple closed curve in G .

More precisely:

Theorem (Residue Theorem). *Let G be an open set, E a discrete subset of G , and γ a null-homotopic piecewise smooth closed curve in G which doesn't intersect E . Then $\{a \in E | n(\gamma; a) \neq 0\}$ is finite and*

$$\frac{1}{2\pi i} \int_{\gamma} f(z) dz = \sum_{a \in E} n(\gamma; a) \text{Res}(f; a).$$

for all functions f which are analytic in $G \setminus E$.

Proof. Let $F : [0, 1] \times [0, 1] \rightarrow G$ be a homotopy from γ to a constant. The image K of F is compact, so has a finite intersection with E . If $a \in E$ does not belong to this intersection, the homotopy takes place in the complement of a , showing that the $n(\gamma; a) = 0$.

Let a_1, \dots, a_m be the points in the intersection, and let g_i be the singular part of f at a_i . The $f - \sum g_i$ is analytic on K (after removing the removable singularities at the a_i), so $\int_{\gamma} f - \sum g_i = 0$. The result follows easily. \square

We will proceed to two easy but useful consequences of the residue theorem, the argument principle and Rouché's theorem. First, however, we consider the application of the residue theorem to the computation of definite integrals.

Residue integrals.

First, how to practically compute residues. For essential singularities it can be very difficult, but for poles it is usually easy. If f has a simple pole at a , then $f(z) = g(z)/(z-a)$ with g analytic at a and $\text{Res}(f; a) = g(a)$ (and thus Cauchy's theorem may be viewed as the special case of the residue theorem where f has a simple pole). If f has a double pole at a , then

$$f(z) = g(z)/(z-a)^2 = \frac{g(a)}{(z-a)^2} + \frac{g'(a)}{z-a} + \dots,$$

so $\text{Res}(f; a) = g'(a)$. Similarly, if f has a pole of order m with $f(z) = g(z)/(z - a)^m$, we expand g as a Taylor series around a and can read off $\text{Res}(f; a)$. (It is $\frac{1}{(m-1)!}g^{(m-1)}(a)$, but it is easier to remember the procedure than the formula.)

Example 1: evaluate
$$\int_{-\infty}^{\infty} \frac{1}{(x^2 + a^2)(x^2 + b^2)} dx, \quad 0 < a < b.$$

Use a semicircle centered at the origin in upper half plane and let radius tend to infinity to get the answer of $\pi/[ab(a + b)]$. Same technique works for all rational functions of degree ≤ -2 .

Example 2: evaluate
$$\int_0^{2\pi} \sin^4(x) dx.$$

Write as $\int_{\gamma} \frac{1}{iz} \left(\frac{z+1/z}{2i}\right)^4 dz$, with γ the unit circle, to get the answer $3\pi/4$. This approach applies to rational functions in \sin and \cos .

Example 3: evaluate
$$\int_0^{\infty} \frac{1}{t^{1/2} + t^{3/2}} dt.$$

Use a branch of the $z^{-1/2}$ with the *positive* real axis as the branch cut, and integrate around this cut and over a large circle centered at the origin (avoid origin with a small circle). The answer is π . This approach applies to integrals of the form $\int_0^{\infty} t^{\alpha} r(t) dt$ where $\alpha \in (0, 1)$ and $r(t)$ is a rational function of degree ≤ -2 with at worst a simple pole at 0.

Example 4: evaluate
$$\int_0^{\infty} \frac{\sin x}{x} dx$$

We will evaluate

$$\lim_{\substack{X \rightarrow +\infty \\ \delta \downarrow 0}} \int_{-X}^{-\delta} + \int_{\delta}^X \frac{e^{iz}}{z} dz.$$

The imaginary part of the quantity inside the limit is $2 \int_{\delta}^X (\sin x)/x dx$. Note: it is essential that when taking the limit we excise a symmetric interval about zero. (The ordinary integral doesn't exist: e^{ix}/x has a non-integrable singularity near the origin, even though its imaginary part, $(\sin x)/x$ doesn't.) On the other hand, it is *not* necessary that we cut off the integrand at symmetric points $-X$ and X (we could have used $-X_1$ and X_2 and let them go independently to zero).

We use a somewhat fancy path: starting at $-X$ we follow the x -axis to $-\delta$, then a semicircle in the upper-half plane to δ , then along the x -axis to X , then along a vertical segment to $X + iX$, then back along a horizontal segment to $-X + iX$, and then down a vertical segment back to $-X$. The integrand e^z/z has no poles inside this curve, so its contour integral is 0. It is easy to see that the integral is bounded by $1/X$ on each of the vertical segments and by $2e^{-X}$ on the horizontal segment. On the other hand,

$e^{iz}/z = 1/z + \text{analytic}$ and the integral of $1/z$ over the semicircle (integrated clockwise) is easily seen to be $-\pi i$. We conclude that the limit above equals πi , whence

$$\int_0^\infty \frac{\sin x}{x} dx = \frac{\pi}{2}.$$

Example 5: evaluate
$$\sum_{n=-\infty}^{\infty} \frac{1}{(n-a)^2}, \quad a \in \mathbb{R} \setminus \mathbb{Z}.$$

Use a rectangular path with opposite corners $\pm(n+1/2+ni)$, n a positive integer and evaluate $\int_\gamma (z-a)^{-2} \cot \pi z dz$, $a \notin \mathbb{Z}$. The poles of the integrand are at a and the integers and the residue is $-\pi/\sin^2(\pi a)$ and $1/[\pi(n-a)^2]$. Since $\sin x + iy = \sin x \cosh y + i \cos x \sinh y$ and $\cos x + iy = \cos x \cosh y - i \sin x \sinh y$, we get

$$|\cot x + iy|^2 = \frac{\cos^2 x + \sinh^2 y}{\sin^2 x + \sinh^2 y},$$

so the $|\cot|$ equals 1 on the vertical segments and, if n is large, is bounded by 2 on the horizontal segments. It follows easily that the integral goes to 0 with n , so

$$\sum_{n=-\infty}^{\infty} \frac{1}{(n-a)^2} = \frac{\pi^2}{\sin^2 \pi a}.$$

To state the argument principle we define:

Definition. A complex function on an open set G is called *meromorphic* if it is analytic on G except for a set of poles.

Note that the pole set of a meromorphic function is discrete (but may be infinite). A meromorphic function is continuous when viewed as taking values in \mathbb{C}_∞ .

Suppose f is meromorphic at a (i.e., on a neighborhood of a). Then $\text{Res}(f/f'; a) = \text{ord}_a(f)$. Indeed, $f(z) = (z-a)^m g(z)$ with $m = \text{ord}_a(f)$ and $g(a) \neq 0$, and $f'/f = m/(z-a) + g'/g$. The residue theorem then immediately gives

Theorem (Argument Principle). *Let G be an open set, f a meromorphic function on G , and γ a null-homotopic piecewise smooth closed curve in G which doesn't intersect either the zero set Z or the pole set P of f . Then*

$$\frac{1}{2\pi i} \int_\gamma \frac{f'}{f} = \sum_a n(\gamma; a) \text{ord}_a(f)$$

where the sum is over $Z \cup P$ and contains only finitely many terms.

Note that the left hand side is nothing other than $n(f \circ \gamma; 0)$, which is behind the name (how much does the argument change as we traverse this curve). We saw this theorem previously in the case where f was analytic, in which case the integral counts the zeros of f . For meromorphic f we count the poles as negative zeros.

Our final application of the residue theorem is Rouché's theorem.

Theorem (Rouché). *Let G be an open set, f and g meromorphic functions on G , and γ a null-homotopic piecewise smooth closed curve in G which doesn't intersect $Z_f \cup P_f \cup Z_g \cup P_g$. If*

$$|f(z) + g(z)| < |f(z)| + |g(z)|, \quad z \in \gamma,$$

then

$$\sum_{a \in Z_f \cup P_f} n(\gamma; a) \operatorname{ord}_a(f) = \sum_{a \in Z_g \cup P_g} n(\gamma; a) \operatorname{ord}_a(g).$$

Proof. The hypothesis means that on γ the quotient $f(z)/g(z)$ is not a non-negative real number. Let l be a branch of the logarithm on $\mathbb{C} \setminus [0, \infty)$. Then $l(f/g)$ is a primitive of $(f/g)'/(f/g) = f'/f - g'/g$ which is analytic in a neighborhood of γ . Hence, $\int_{\gamma} f'/f = \int_{\gamma} g'/g$ and the theorem follows from the argument principle. \square

Corollary (Classical statement of Rouché's theorem). *Suppose f and ϕ are analytic on a simply-connected region G and that γ is a simple closed piecewise smooth curve in G . If $|\phi| < |f|$ on γ , then $f + \phi$ has the same number of zeros as f in the region bounded by γ .*

Proof. $|(f + \phi) + (-f)| = |\phi| < |f| \leq |f + \phi| + |-f|$. \square

Example: If $\lambda > 1$, then the equation $z \exp(\lambda - z) = 1$ has exactly one solution in the unit disk (take $f = z \exp(\lambda - z)$, $\phi = -1$).

V. FURTHER RESULTS ON ANALYTIC FUNCTIONS

The theorems of Weierstrass, Hurwitz, and Montel. Consider the space of continuous functions on a region in the complex plane, endowed with the topology of uniform convergence on compact subsets. (In fact this is a metric space topology: we can define a metric such that a sequence of continuous functions converge to another continuous function in this metric, if and only if the sequence converges uniformly on compact subsets.) Weierstrass's theorems states that the space of analytic functions on the region is a closed subspace and differentiation acts continuously on it. Montel's theorem identifies the precompact subsets of the space of analytic functions.

Theorem (Weierstrass). *Let $G \subset \mathbb{C}$ be open, $f_n : G \rightarrow \mathbb{C}$, $n = 1, 2, \dots$, analytic, $f : G \rightarrow \mathbb{C}$. If $f_n \rightarrow f$ uniformly on compact subsets of G , then f is analytic. Moreover f'_n converges to f' uniformly on compact subsets of G .*

Proof. If T is a triangular path,

$$\int_T f = \lim_{n \rightarrow \infty} \int_T f_n = 0,$$

so f is analytic by Morera's theorem.

Let γ be the boundary of a disk around z contained in G .

$$\lim f'_n(z) = \lim \frac{1}{2\pi i} \int_{\gamma} \frac{f'_n(w)}{(w-z)^2} dz = \frac{1}{2\pi i} \int_{\gamma} \frac{f'(w)}{(w-z)^2} dz = f'(z). \quad \square$$

Combining Weierstrass's Theorem and the Argument Principle we get:

Theorem (Hurwitz). *Let G be a domain in \mathbb{C} , f_n, f analytic functions on G with $f_n \rightarrow f$ uniformly on compact subsets. If each of the f_n is nowhere-vanishing, then f is either nowhere-vanishing or identically zero.*

Proof. If f is not identically zero, but $f(a) = 0$, we can choose $r > 0$ so that $\bar{B}(a, r) \subset G$, and a is the only zero of f on this disk. Letting γ be the boundary of the disk, the argument principle then tells us that $\int_{\gamma} f'_n/f_n = 0$, but $\int_{\gamma} f'/f = 2\pi i \operatorname{ord}_a(f) \neq 0$. But, in view of Weierstrass's Theorem the hypothesis tells us that f'_n/f_n converges to f'/f uniformly on γ , so we have a contradiction. \square

Corollary. *Let G be a domain in \mathbb{C} , f_n, f analytic functions on G with $f_n \rightarrow f$ uniformly on compact subsets. If each of the f_n is injective, then f is either injective or identically constant.*

Proof. If f is not identically constant, but there exists distinct a, b with $f(a) = f(b)$, then we can choose $r > 0$ so that $B(a, r) \cup B(b, r) \subset G$, and $B(a, r) \cap B(b, r) = \emptyset$. Now $f_n(z) - f(a)$ converges to $f(z) - f(a)$ uniformly on compact subsets of $B(a, r)$ and the latter function has a zero but is not identically zero there, so for all sufficiently large n , $f_n(z) = f(a)$ for some $z \in B(a, r)$, and, by the same argument, for some $z \in B(b, r)$. This contradicts the injectivity of the f_n . \square

Theorem (Montel). *Let $G \subset \mathbb{C}$ be open, \mathcal{F} a family of analytic functions on G . Suppose that the functions f_1, f_2, \dots are uniformly bounded on each compact subset of G . Then there exists a subsequence which converges uniformly on compact subsets.*

Proof. Fix $a \in G$ and let $d(a) = \operatorname{dist}(a, \partial G)/2$. First we show that we can find a subsequence that converges uniformly on $B(a, d(a)/2)$. Now

$$f_m(z) = \sum_{n=0}^{\infty} c_n(f_m)(z-a)^n, \quad z \in \bar{B}(z, d(a)).$$

where $c_n(f) = f^{(n)}(a)/n!$. By the Cauchy estimates

$$|c_n(f_m)| \leq M/d(a)^n$$

where M is the uniform bound for the f_m on $\bar{B}(a, d(a))$.

Since $c_1(f_m)$ is uniformly bounded in \mathbb{C} , we can find a subsequence $f_{m^1(j)}$ for which $c_1(f_{m^1(j)})$ converges as $j \rightarrow \infty$. Then we can find a further subsequence so that $c_2(f_{m^2(j)})$ converges, as well, etc. For the diagonal sequence $g_j = f_{m^j(j)}$ $c_n(g_j)$ converges for all n .

Claim: g_j converges uniformly on $B(a, d(a)/2)$. To prove this, we must show that for any $\epsilon > 0$

$$|g_j - g_k| \leq \epsilon \quad \text{on } B(a, d(a)/2).$$

Now, for any $N \in \mathbb{N}$, $z \in B(a, d(a)/2)$,

$$\begin{aligned} |g_j(z) - g_k(z)| &\leq \sum_{n=1}^N |c_n(g_j) - c_n(g_k)| |z - a|^n + \sum_{n=N+1}^{\infty} \frac{2M}{d(a)^n} |z - a|^n \\ &\leq \sum_{n=1}^N |c_n(g_j) - c_n(g_k)| [d(a)/2]^n + \frac{2M}{2^N}. \end{aligned}$$

Given $\epsilon > 0$ we can choose N large enough that $\frac{2M}{2^N}$ and then choose j and k large enough that $|c_n(g_j) - c_n(g_k)| \leq \epsilon/[2(N+1)[d(a)/2]^n]$ for $0 \leq n \leq N$. This proves the claim.

Fix $a \in G$ and let $d(a) = \text{dist}(a, \partial G)/2$. First we show that we can find a subsequence that converges uniformly on $B(a, d(a)/2)$. Now

$$f_m(z) = \sum_{n=0}^{\infty} \frac{f_m^{(n)}(a)}{n!} (z - a)^n, \quad z \in \bar{B}(a, d(a)).$$

By the Cauchy estimates

$$|f_m^{(n)}(a)| \leq Mn!/d(a)^n$$

where M is the uniform bound for the f_m on $\bar{B}(a, d(a))$. In particular, $\sup_m |f_m^{(n)}(a)| < \infty$ for each n .

Thus we can find a subsequence $f_{m^0(j)}$ for which $f_{m^0(j)}(a)$ converges as $j \rightarrow \infty$. Then we can find a further subsequence so that $f'_{m^2(j)}(a)$ converges as well, etc. For the diagonal sequence $g_j = f_{m^j(j)}(a)$ converges for all n .

Claim: g_j converges uniformly on $B(a, d(a)/2)$. To prove this, we must show that for any $\epsilon > 0$

$$|g_j - g_k| \leq \epsilon \quad \text{on } B(a, d(a)/2).$$

Now, for any $N \in \mathbb{N}$, $z \in B(a, d(a)/2)$,

$$\begin{aligned} |g_j(z) - g_k(z)| &\leq \sum_{n=0}^N |g_j^{(n)}(a) - g_k^{(n)}(a)| |z - a|^n / n! + \sum_{n=N+1}^{\infty} \frac{2M}{d(a)^n} |z - a|^n \\ &\leq \sum_{n=1}^N |g_j^{(n)}(a) - g_k^{(n)}(a)| |z - a|^n / n! + \frac{2M}{2^N}. \end{aligned}$$

Given $\epsilon > 0$ we can choose N large enough that $2M/2^N \leq \epsilon/2$ and can then choose j and k large enough that $|g_j^{(n)}(a) - g_k^{(n)}(a)| \leq n!\epsilon/\{2(N+1)[d(a)/2]^n\}$ for $0 \leq n \leq N$. This proves the claim and shows that we can find a subsequence which converges uniformly on $B(a, d(a)/2)$.

Now we may choose a countable sequence of points $a_i \in G$ such that $G = \bigcup_{i=1}^{\infty} B(a_i, d(a_i)/2)$ (e.g., all points in G with rational real and imaginary parts). Given the sequence of functions f_m uniformly bounded on compact sets, we may find a subsequence $f_{n^1(j)}$ which converges uniformly on $B(a_1, d(a_1)/2)$. Then we may take a subsequence of that sequence which converges uniformly on $B(a_2, d(a_2)/2)$, etc. The diagonal subsequence h_j converges uniformly on each $B(a_i, d(a_i)/2)$.

We complete the proof by showing that h_j converges uniformly on all compact subsets. Indeed this is immediate, because any compact subset of G is contained in a finite union of the $B(a_i, d(a_i)/2)$. \square

Schwarz's Lemma. Schwarz's lemma is an easy, but very useful, consequence of the maximum modulus principle. The most difficult part is to remember that the German analyst Herman Schwarz (actually Karl Herman Amandus Schwarz, 1843-1921), inventor of the "Schwarz inequality," "Schwarz lemma," and "Schwarz alternating method," spells his name without a "t", while the French analyst, Laurent Schwartz, 1915–, inventor of the theory of distributions, uses a "t".

Suppose that an analytic map f maps a disk of radius ϵ about a into the disk of radius δ about $f(a)$. We can use this to estimate the derivative $f'(a)$ and the stretching $|f(b) - f(a)|/|b - a|$. To keep the statement simple, we translate in the domain and range space so that $a = f(a) = 0$ and dilate in the domain and range so that $\epsilon = \delta = 1$. We then get:

Theorem (Schwarz's Lemma). *Suppose f maps the open unit disk into itself leaving the origin fixed. Then $|f(z)| \leq |z|$ for all z in the disk, and if equality holds for any nonzero point, then $f(z) = cz$ for some c of modulus 1. Also $|f'(0)| \leq 1$ and if equality holds, then $f(z) = cz$ for some c of modulus 1.*

Proof. The function $g(z) = f(z)/z$ has a removable singularity at the origin, and extends analytically to the disk with $g(0) = f'(0)$. It satisfies $|g(z)| \leq 1/r$ on the circle of radius $r < 1$, so by the maximum modulus theorem it satisfies this condition on the disk of radius r . Letting r tend to one gives the result. Equality implies g is constant. \square

From a homework exercise (§III .3, no. 10), we know that the most general Möbius transformations of the open unit disk D into itself is given by

$$z \mapsto c \frac{z - a}{1 - \bar{a}z}$$

with a in the disk and $|c| = 1$. We can use Schwarz's lemma to show that this is most general one-to-one analytic map of the disk onto itself.

First consider the special case $f(0) = 0$. Then Schwarz shows that $|f'(0)| \leq 1$ with equality if and only if $f(z) = cz$ some $|c| = 1$. Applying Schwarz to f^{-1} gives $|1/f'(0)| = |(f^{-1})'(0)| \leq 1$. Thus equality does indeed hold.

Returning to the general case, say $f(a) = 0$. Then we can apply the special case above to $f \circ \phi_{-a}$ to get $f(\phi_{-a}(z)) = cz$ or $f(z) = c\phi_a(z)$.

The Riemann Mapping Theorem. We begin by deriving some easy consequences of simple-connectivity. Recall that every analytic function on a simply-connected region admits a primitive. It follows that if f is a nowhere-vanishing function on a simply-connected domain, then there exists an analytic function F on that domain such that $f = \exp(F)$. Indeed, let g be primitive of f'/f . It follows immediately that $f \exp(-F)$ is constant, and so adjusting g by an additive constant, $f = \exp(F)$. Another consequence is the *square-root property*: if f is a nowhere-vanishing analytic function on the domain, then there exists an analytic function g with $[g(z)]^2 = f(z)$. Indeed, we can just take $g = \exp(F/2)$. For future use we remark that if there exists an analytic isomorphism of a region G_1 onto a region G_2 , then G_1 has the square-root property if and only if G_2 does.

We now turn to a lemma of Koebe.

Lemma. *Let $D = B(0, 1)$ and let $G \subsetneq D$ be a region containing the origin and having the square root property. Then there exists an injective analytic map $r : G \rightarrow D$ such that $r(0) = 0$, $|r(z)| > |z|$, $z \in G \setminus \{0\}$.*

Proof. Let $a \in D \setminus G$ and let b be one of the square roots of $-a$. Define

$$q(z) = \phi_{-a}([\phi_{-b}(z)]^2)$$

(where $\phi_\alpha(z) := (z - \alpha)/(1 - \bar{\alpha}z)$). Clearly $q(0) = 0$, and q is not just multiplication by a constant (in fact, it is not 1-1 on D), so Schwarz's lemma implies

$$|q(z)| < |z|, \quad 0 \neq z \in D.$$

Next, note that ϕ_a is an analytic map of G into D which never vanishes, so admits an analytic square root $g(z)$, which we may determine uniquely by insisting that $g(0) = b$. Defining

$$r = \phi_b \circ g,$$

we have $r(0) = 0$, and $q(r(z)) \equiv z$. Thus if $0 \neq z \in G$, $r(z) \neq 0$ and $|z| = |q(r(z))| < |r(z)|$. \square

Two regions in the complex plane are called conformally equivalent if they are analytically isomorphic, that is, if there exists a one-to-one analytic mapping of the first domain onto the second (whose inverse is then automatically analytic).

Riemann Mapping Theorem. *Let $G \subsetneq \mathbb{C}$ be simply-connected. Then G is conformally equivalent to the open unit disk.*

Proof. The only consequence of simple-connectivity that we shall use in the proof is that G has the square-root property.

The structure of the proof is as follows: first we use the hypotheses to exhibit an injective analytic map of h_0 of G onto a domain G_0 which satisfies the hypotheses of Koebe's lemma. Then we use an extremal problem to define an injective analytic map f of G_0 onto a domain

G_1 contained in the disk. If G_1 fills the entire disk we are done; otherwise, G_1 satisfies the hypotheses of Koebe's lemma and we use the lemma to contradict the extremality of f .

Let $a \in \mathbb{C} \setminus G$ and let g be an analytic square root of $z - a$. It is easy to see that g is injective and that if $w \in g(G)$ then $-w \notin g(G)$. Pick $w_0 \in g(G)$ and then $r > 0$ such that $B(w_0, r) \subset g(G)$ (Open Mapping Theorem). Therefore $B(-w_0, r) \cap g(G) = \emptyset$. Set $h(z) = [g(z) + w_0]^{-1}$ so that $h : G \rightarrow \bar{B}(0, 1/r)$ is analytic and injective. Then set $h_0(z) = r[h(z) - h(z_0)]/3$ where $z_0 \in G$. This map is an analytic isomorphism of G onto a connected open set $G_0 \subsetneq D$ containing 0.

Let \mathcal{F} be the set of all injective analytic functions $f : G_0 \rightarrow D$ with $f(0) = 0$. Fix $0 \neq w_0 \in G_0$, and let $\alpha = \sup_{f \in \mathcal{F}} |f(w_0)|$. Note $\alpha \in (0, 1]$. We claim that the supremum is achieved. Indeed, let $f_n \in \mathcal{F}$, $|f_n(w_0)| \rightarrow \alpha$. Since the f_n are uniformly bounded (by 1), Montel's theorem assures us that there is a subsequence f_{n_k} which converges uniformly on compact subsets to some analytic function f on G_0 . Obviously $f(0) = 0$ and $|f(w_0)| = \alpha > 0$, so f is not constant, and since the f_n are injective, so is f . Clearly $|f| \leq 1$ on G_0 , and by the maximum modulus principle, $|f| < 1$. Thus $f \in \mathcal{F}$ and achieves the supremum.

We complete the theorem by showing that f is an isomorphism of G_0 onto D (so $f \circ h_0$ is the desired analytic isomorphism of G onto D). We already know that f is analytic and injective, so we need only show that f maps onto D . If, to the contrary, $G_1 := f(G_0) \subsetneq D$ then (since G_1 inherits the square-root property from G_0 which inherits it from G) Koebe's lemma gives an injective analytic function $r : G_1 \rightarrow D$ with $r(0) = 0$, $|r(z)| > |z|$ for $z \neq 0$. Then $r \circ f \in \mathcal{F}$, but $|r(f(w_0))| > |f(w_0)|$, a contradiction. \square

We now note that the square-root property is equivalent to simple-connectivity. Indeed we have seen that simple-connectivity implies the square-root property. On the other hand, if G satisfies the square root property, then either $G = \mathbb{C}$, which is simply-connected, or G is isomorphic to the disk (by the proof of the Riemann mapping theorem above), and so is simply-connected.

If $G \subsetneq \mathbb{C}$ is a simply-connected region and $z_0 \in G$ is arbitrary we may take any analytic isomorphism of G onto the disk and follow it with a suitably chosen Möbius transformation ($z \mapsto c(a - z)/(1 - \bar{a}z)$, $|a| < 1$, $|c| = 1$, to obtain an analytic isomorphism f of G onto D satisfying

$$f(z_0) = 0, \quad f'(z_0) > 0.$$

If f_1 and f_2 were two such analytic isomorphisms, then $g = f_2 \circ f_1^{-1}$ is an analytic isomorphism of the disk onto itself with $g(0) = 0$, $g'(0) > 0$, and it easily follows that g is the identity and $f_1 = f_2$.

Complements on Conformal Mapping. There are many interesting questions in conformal mapping which we will not have time to investigate, but I will quickly state some key results.

Extension to boundary. If the boundary of a simply-connected region is a simple closed curve, then it can be proved that a conformal map of the region onto the open disk extends to a topological homeomorphism of the closure of the region on to the closed disk.

Constructive conformal mapping. Conformal is used to obtain explicit solutions to problems involving analytic and harmonic functions. For example, Joukowski used the conformal mapping $f \circ g$ where

$$f(z) = 2\frac{1+z}{1-z}, \quad g(z) = \left(\frac{z-1}{z+1}\right)^2,$$

applied to a circle passing through -1 and containing 1 in its interior. The resulting region, called a Joukowski airfoil, looks like a cross-section of a wing, and by adjusting the circle, one can adjust the exact shape of the airfoil. Using the explicit conformal map to the disk and explicit formulas for the solution of Laplace's equation on (the exterior of) a disk (similar to those that will be derived in the next section, one can calculate things like the drag and lift of the airfoil, and so this was a useful approach to airfoil design.

An important class of regions are the polygons. The Schwarz–Christoffel formula is an explicit formula for the conformal map of a given polygon onto the disk. The formula involves a few complex parameters which are determined by the vertices of the polygon. Given the vertices, one cannot usually compute the parameters analytically, but they are not difficult to compute numerically (with a computer).

Multiply connected regions. For lack of time we will not study conformal mapping of multiply connected regions, but we just state, without proof, the result for doubly connected regions (regions whose complement in the Riemann sphere consists of two connected components, i.e., roughly regions in the plane with one hole). It can be shown (with basically the tools at our disposal) that any such region is conformally equivalent to the annulus $1 < |z| < R$ for some $R > 1$. The value of R is uniquely determined. In particular, two such annuli with different values of R_1 are *not* conformally equivalent.

VI. HARMONIC FUNCTIONS

A real-valued C^2 function on an open subset of \mathbb{C} is called *harmonic* if its Laplacian vanishes. We have seen previously that the real part of an analytic function is harmonic and that if the domain is simply-connected every real-valued harmonic function u admits a harmonic conjugate, i.e., a real-valued function v , for which $u+iv$ is analytic. In particular, we see that every harmonic function is C^∞ .

Suppose u is harmonic on G , $a \in G$. If $r > 0$ is such that $\bar{B}(a, r)$, then we can choose a disk D with $\bar{B}(a, r) \subset D \subset G$, and since D is simply-connected $u = \operatorname{Re} f$ on D for some analytic function f on D . Cauchy's integral formula then gives

$$f(a) = \frac{1}{2\pi i} \int_{|z-a|=r} \frac{f(z)}{z-a} dz = \frac{1}{2\pi} \int_0^{2\pi} f(a + re^{i\theta}) d\theta.$$

Taking real parts we get:

Mean Value Theorem. *If $u : G \rightarrow \mathbb{R}$ is harmonic and $\bar{B}(a, r) \subset G$, then*

$$(MVP) \quad u(a) = \frac{1}{2\pi} \int_0^{2\pi} u(a + re^{i\theta}) d\theta.$$

A harmonic function satisfies the *Maximum Principle*: it does not assume its maximum on a region unless it is constant there. Indeed the conclusion is true for any continuous function with the mean-value property (MVP), or even just satisfying the inequality $u(a) \leq (2\pi)^{-1} \int_0^{2\pi} u(a + re^{i\theta}) d\theta$ (since this implies that the set of points where the maximum is achieved is open, and it is obviously closed). The reverse inequality similarly shows that a harmonic function satisfies the Minimum Principle.

Given a bounded domain G and a function $f : \partial G \rightarrow \mathbb{R}$, the *Dirichlet problem* consists of finding a function $u : \bar{G} \rightarrow \mathbb{R}$ such that u is harmonic on G , continuous on \bar{G} , and coincident with f on ∂G . An important consequence of the maximum and minimum principles is uniqueness for the Dirichlet problem. If u_1 and u_2 are both harmonic on G , continuous on \bar{G} , and agree on ∂G , then $u_1 - u_2$ is harmonic and vanishes on the boundary, but it takes its maximum and minimum on the boundary, so it is identically zero, i.e., $u_1 = u_2$.

When applying the maximum principle to functions which are not known to extend continuously to the boundary this easy-to-prove consequence is useful: If u satisfies the maximum principle on a bounded open set and there is a number K such that $\limsup_{z \rightarrow z_0} u(z) \leq K$, then $u \leq K$ on G . [Proof: Let $M = \sup_{z \in G} u(z)$ and take $z_n \in G$ with $u(z_n) \rightarrow M$. Pass to a subsequence with $z_n \rightarrow z_0 \in \bar{G}$. If $z_0 \in G$, then u takes its maximum there, so is constant and obviously $M \leq K$. Otherwise $z_0 \in \partial G$, so $K \geq \limsup_{z \rightarrow z_0} u(z) \geq \lim_n u(z_n) = M$.]

The Poisson kernel. For $0 \leq r < 1$, $\theta \in \mathbb{R}$ let

$$P_r(\theta) = \operatorname{Re} \left(\frac{1 + re^{i\theta}}{1 - re^{i\theta}} \right) = \frac{1 - r^2}{1 - 2r \cos \theta + r^2},$$

define the Poisson kernel. Note that

$$P_r(\theta) = \operatorname{Re} \left(1 + 2 \sum_{n=1}^{\infty} r^n e^{in\theta} \right) = \sum_{n=-\infty}^{\infty} r^{|n|} e^{in\theta}.$$

Lemma. For $0 \leq r < 1$, $P_r(\theta)$ is a smooth, positive, even, 2π -periodic function of θ with mean value 1. Moreover if $\theta > 0$, then

$$\lim_{r \uparrow 1} P_r(\theta) = 0, \quad \theta \in \mathbb{R} \setminus 2\pi\mathbb{Z},$$

and if $\delta > 0$, then the convergence is uniform over $\theta \in \mathbb{R}$ such that $|\theta - 2\pi n| \geq \delta$ for all $n \in \mathbb{Z}$.

Proof. Elementary (use the power series expansion for the integral and the formula

$$P_r(\theta) = \frac{1 - r^2}{(1 - r)^2 + 2r(1 - \cos \theta)}$$

for the limits). \square

Now let D be the open unit disk and ∂D its boundary. The next theorem shows how to solve the Dirichlet problem for D using the Poisson kernel:

Theorem. Given $f : \partial D \rightarrow \mathbb{R}$ continuous, define

$$\begin{cases} u(z) = f(z), & z \in \partial D, \\ u(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} P_r(\theta - t) f(e^{it}) dt, & 0 \leq r < 1, \quad \theta \in \mathbb{R}. \end{cases}$$

The u is continuous on \bar{D} and harmonic on D .

Proof.

$$\begin{aligned} u(re^{i\theta}) &= \operatorname{Re} \left\{ \frac{1}{2\pi} \int_0^{2\pi} f(e^{it}) \left[\frac{1 + re^{i(\theta-t)}}{1 - re^{i(\theta-t)}} \right] dt \right\} \\ &= \operatorname{Re} \left\{ \frac{1}{2\pi} \int_0^{2\pi} f(e^{it}) \left[\frac{e^{it} + re^{i\theta}}{e^{it} - re^{i\theta}} \right] dt \right\} \end{aligned}$$

or

$$u(z) = \operatorname{Re} \left\{ \frac{1}{2\pi} \int_0^{2\pi} f(e^{it}) \left[\frac{e^{it} + z}{e^{it} - z} \right] \right\}, \quad z \in D.$$

The integral defines an analytic function of z (since the integrand is continuous in t and z and analytic in z), so u , being the real part of an analytic function, is harmonic.

It remains to show that $\lim_{\substack{z \in D \\ z \rightarrow e^{it_0}}} u(z) = f(e^{it_0})$ for $t_0 \in \mathbb{R}$. Now

$$u(re^{i\theta}) - f(e^{it_0}) = \frac{1}{2\pi} \int_{t_0-\pi}^{t_0+\pi} P_r(\theta - t) [f(e^{it}) - f(e^{it_0})] dt,$$

so it suffices to show that the last integral can be made arbitrarily small by taking θ sufficiently close to t_0 and $r < 1$ sufficiently close to 1. Now given $\epsilon > 0$ we can choose $\delta > 0$ so that $|f(e^{it}) - f(e^{it_0})| \leq \epsilon$ if $|t - t_0| \leq \delta$. Split the interval of integration as

$$\{t : |t - t_0| \leq \pi\} = I_1 \cup I_2 := \{t : |t - t_0| \leq \delta\} \cup \{t : \delta \leq |t - t_0| \leq \pi\}.$$

On I_1 , the integrand is bounded by $P_r(\theta - t)\epsilon$, so the integral is bounded by

$$\frac{\epsilon}{2\pi} \int_{I_1} P_r(\theta - t) dt \leq \frac{\epsilon}{2\pi} \int_0^{2\pi} P_r(\theta - t) dt \leq \epsilon.$$

Now suppose that θ is taken sufficiently near t_0 , namely $|\theta - t_0| \leq \delta/2$. Then for $t \in I_2$,

$$|\theta - t| \geq |t - t_0| - |\theta - t_0| \geq \delta/2, \quad |\theta - t| \leq |t - t_0| + |\theta - t_0| \leq \pi + \delta/2$$

Thus $P_r(\theta - t) \rightarrow 0$ uniformly for $t \in I_2$ as $r \uparrow 1$. Since f is bounded, the entire integrand does so as well. Thus if $|\theta - t_0| \leq \delta/2$ we may take r sufficiently close to 1 that the integral over I_2 is bounded by ϵ . \square

It is now easy to deduce that the mean value property characterizes harmonic functions.

Theorem. *If $u : G \rightarrow \mathbb{R}$ is a continuous function satisfying the mean value property, then it is harmonic.*

Proof. It is enough to show that u is harmonic on each disk with closure in G . From the Poisson kernel construction we can construct a harmonic function with the same boundary values as u on this disk. The difference between this function and u satisfies the mean value property and is zero on the boundary of the disk, so it vanishes on the disk. \square

Corollary. *If a sequence of harmonic functions converges uniformly on compact subsets, then the limit function is harmonic.*

From the expressions

$$P_r(\theta) = \frac{1 - r^2}{(1 + r)^2 - 2r(1 + \cos \theta)} = \frac{1 - r^2}{(1 - r)^2 + 2r(1 - \cos \theta)}$$

we have

$$\frac{1 - r}{1 + r} \leq P_r(\theta) \leq \frac{1 + r}{1 - r}.$$

Now suppose u is harmonic on $B(0, R)$ and non-negative, and $re^{i\theta}$ is an arbitrary point of $B(0, R)$. Choose $\epsilon > 0$ small enough that $r < R - \epsilon$, and set $U(z) = u((R - \epsilon)z)$. U is harmonic on a neighborhood of $\bar{B}(0, 1)$, so

$$u(re^{i\theta}) = U([r/(R - \epsilon)]e^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} P_{r/(R-\epsilon)}(\theta - t)U(e^{it}) dt.$$

Using the non-negativity of u , the mean value property, and the upper bound for P_r , we get

$$u(re^{i\theta}) \leq \frac{1 + r/(R - \epsilon)}{1 - r/(R - \epsilon)} u(0).$$

Letting $\epsilon \downarrow 0$ this gives

$$u(re^{i\theta}) \leq \frac{R + r}{R - r} u(0).$$

We obtain a lower bound in a similar fashion. Of course we can translate the disk so it is centered at an arbitrary point a . Thus we have proven:

Theorem (Harnack's Inequalities). *If u is non-negative and harmonic on $B(a, R)$, then*

$$\frac{R - r}{R + r} u(a) \leq u(a + re^{i\theta}) \leq \frac{R + r}{R - r} u(a)$$

for $0 \leq r < R$.

For example, if u is harmonic and non-negative on the disk of radius R about a then $u(z) \in [u(a)/3, 3u(a)]$ on the disk of radius $R/2$ about a .

Theorem (Harnack). *Let $0 \leq u_1 \leq u_2 \leq \dots$ be a sequence of harmonic functions on a region G . Then either $\lim_n u_n(z) = +\infty$ uniformly on compact subsets of G or there is an harmonic function u with $\lim_n u_n(z) = u(z)$ uniformly on compact subsets of G .*

Proof. For any point $a \in G$ let D be a disk centered at a contained in G . Then there exists a constant $C > 1$ with

$$u_n(z) \leq C u_n(a), \quad u_n(a) \leq C u_n(z), \quad z \in D.$$

Consequently, the numbers $u_n(a)$ remain bounded as $n \rightarrow \infty$ if and only if $u_n(z)$ remains bounded for all $z \in D$. This shows that the set of points where u_n stays bounded is both open and closed, so it is either the empty set or all of G . In the first case, $u_n(z) \rightarrow +\infty$ for all $z \in G$, and the above estimate shows that the convergence is uniform on a covering set of discs, hence on all compact subsets.

In the second case, $\lim_n u_n(a) \in \mathbb{C}$ for all a , and since

$$u_n(z) - u_m(z) \leq C[u_n(a) - u_m(a)], \quad n \geq m, z \in D,$$

the sequence is uniformly Cauchy on D so converges uniformly on D . Again, uniform convergence on compact sets follows. \square

Subharmonic functions and the solution of the Dirichlet problem.

Definition. Let u be a continuous real-valued function on a region G . If

$$u(a) \leq \frac{1}{2\pi} \int_0^{2\pi} u(a + re^{i\theta}) d\theta$$

whenever $\bar{B}(a, r) \subset G$, we say that u is *subharmonic*.

As mentioned earlier, a subharmonic function satisfies the maximum principle. Other simple properties are:

- harmonic functions are subharmonic
- the sum of two subharmonic functions is subharmonic
- the pointwise maximum of two subharmonic functions is subharmonic

The next lemma states that subharmonicity is a local property:

Lemma. *Suppose that u is subharmonic in a neighborhood of each point of a region G . Then u is subharmonic on G .*

Proof. If $\bar{D} = \bar{B}(a, r) \subset G$, let \tilde{u} be the continuous function on \bar{D} which agrees with u on ∂D and is harmonic on D . From the Poisson kernel representation, we have

$$\tilde{u}(a) = \frac{1}{2\pi} \int_0^{2\pi} u(a + re^{i\theta}) d\theta.$$

Now $u - \tilde{u}$ is subharmonic in a neighborhood of each point of D , and it follows by the usual connectedness argument that the points where it achieves its maximum on D is either empty or all of D . Hence its maximum on \bar{D} is achieved on the boundary, so is 0. This shows that $u(a) \leq \tilde{u}(a)$ as desired. \square

One more property we shall use:

Lemma. Suppose that u is subharmonic in a region G and $\bar{D} = \bar{B}(a, r) \subset G$. Define \tilde{u} on \bar{D} as the continuous function on \bar{D} which agrees with u on ∂D and is harmonic on D , and set $\tilde{u} = u$ on $G \setminus \bar{D}$. Then \tilde{u} is subharmonic on G .

Proof. Clearly it is continuous and subharmonic in D and $G \setminus \bar{D}$. So it suffices to show that for each point a of ∂D , \tilde{u} is bounded by its mean on any circle around a . Now $u - \tilde{u}$ is subharmonic on D and 0 on ∂D , so $u \leq \tilde{u}$ on D , and therefore on G . Thus

$$\tilde{u}(a) = u(a) \leq \frac{1}{2\pi} \int_0^{2\pi} u(a + re^{i\theta}) d\theta \leq \frac{1}{2\pi} \int_0^{2\pi} \tilde{u}(a + re^{i\theta}) d\theta.$$

□

Note that

Let G be a bounded region. If u is a harmonic function on G continuous up to the boundary, and v is any subharmonic function that is less than u on the boundary, then the maximum principle implies that $v \leq u$ in G . Hence it is reasonable to try to solve the Dirichlet problem by seeking the largest subharmonic function that doesn't exceed the given boundary values. This is the approach of O. Perron.

Let f be a bounded (at this point not necessarily continuous) real-valued function on ∂G . Define the Perron family $\mathcal{P}(f, G)$ as the set of subharmonic functions v on G for which

$$(\star) \quad \limsup_{z \rightarrow a} v(z) \leq f(a) \quad \text{for all } a \in \partial G.$$

Note that $\mathcal{P}(f, G)$ contains all constants not exceeding $\min f$, so it is not empty.

From the maximum principle we obtain:

Lemma. If $v \in \mathcal{P}(f, G)$ then $v \leq \sup f$ on G .

Now define the *Perron function*

$$u(z) = \begin{cases} \sup_{v \in \mathcal{P}(f, G)} v(z), & z \in G, \\ f(z), & z \in \partial G. \end{cases}$$

We will show that (1) the Perron function is harmonic on G , and (2) under mild restrictions on G , it extends continuously to \bar{G} with value f on ∂G . Thus we will have exhibited a solution to the Dirichlet problem.

Proof that the Perron function is harmonic. Let D be any disk with closure in G and let a be any point in the disk. Choose functions $v_n \in \mathcal{P}(f, G)$ with $v_n(a) \rightarrow u(a)$. Let $V_n = \max(v_1, \dots, v_n)$, and then define \tilde{V}_n to be V_n off D and to be harmonic on D and continuous on \bar{D} . We know that V_n and \tilde{V}_n are subharmonic, and clearly the V_n and thus the \tilde{V}_n are non-decreasing. All belong to $\mathcal{P}(f, G)$ so are bounded by u , but $v_n \leq V_n \leq \tilde{V}_n$, so $\tilde{V}_n(a) \uparrow u(a)$. By Harnack's Theorem, \tilde{V}_n converges to an harmonic function V on D .

We know that $V(a) = u(a)$. We will now show that if b is any other point of D then $V(b) = u(b)$. This will show that u is harmonic in D as desired.

Choose functions $\phi_n \in \mathcal{P}(f, G)$ with $\phi_n(b) \rightarrow u(b)$, and set $w_n = \max(\phi_n, v_n) \in \mathcal{P}(f, G)$. Construct W_n and \tilde{W}_n from w_n in analogy with V_n and \tilde{V}_n . Then $w_n \geq v_n$, $W_n \geq V_n$, $\tilde{W}_n \geq \tilde{V}_n$, and W_n converges to a harmonic function $W \geq V$ with $W(b) = u(b)$. Also $W(a) = u(a) = V(a)$, so $V - W$ is a non-positive harmonic function on D which achieves its maximum at a . Therefore $V \equiv W$ on D , so indeed $V(b) = u(b)$. \square

We now turn to the continuity of the Perron solution at the boundary.

Definition. Let G be a bounded region and $z_0 \in \partial G$. A continuous function ϕ on \bar{G} which is harmonic on G is called a *barrier function* for G at z_0 if it is positive on $\bar{G} \setminus \{z_0\}$ and zero at z_0 .

For example, suppose that there is a line through z_0 such that $\bar{G} \setminus \{z_0\}$ lies inside one of the open half-planes determined by the line. We may write the half-plane as $\text{Im}[(z - z_0)/b] > 0$ for some $b \in \mathbb{C}$, and so $\text{Im}[(z - z_0)/b]$ is a barrier function for G at z_0 .

Next consider the function $\sqrt{1 - 1/z}$ where we use the principal branch of the square root, defined on $\mathbb{C} \setminus (-\infty, 0]$. This function is analytic on $\mathbb{C} \setminus [0, 1]$ and extends continuously to 1 with value 0, and its real part is everywhere positive on $\mathbb{C} \setminus [0, 1]$. Thus if $z_0 = 1$ belongs to ∂G , and is the only point of intersection of the interval $[0, 1]$ with \bar{G} , we can use $\text{Re} \sqrt{1 - 1/z}$ as a barrier function for G at z_0 . Similarly, if z_0 is any boundary point for which there exists a line segment which intersects \bar{G} only at z_0 , then there is a barrier function for G at z_0 . Thus barrier functions exist at all boundary points of smooth domains, polygonal domains, even on a disk minus a segment, and on many other domains. A standard example of a domain without a barrier at some boundary point is a punctured disk.

Theorem. Let G be a bounded domain, f a bounded function on ∂G , and u the corresponding Perron function. If z_0 is a point in ∂G possessing a barrier function and f is continuous at z_0 , then $\lim_{z \rightarrow z_0} u(z) = f(z_0)$.

Proof. First we show that for any $\epsilon > 0$, there is a harmonic function w on G , continuous on \bar{G} , which satisfies $w > f$ on ∂G and $w(z_0) = f(z_0) + \epsilon$. Indeed, let D be a disk about z_0 such that $|f(z) - f(z_0)| < \epsilon$ for $z \in \partial G \cap D$. Let M denote the maximum of $|f|$ on ∂G and m denote to the minimum of the barrier function ϕ on $G \setminus D$, so $m > 0$. Set

$$w(z) = \frac{\phi(z)}{m} [M - f(z_0)] + f(z_0) + \epsilon = \left[\frac{\phi(z)}{m} - 1 \right] [M - f(z_0)] + M + \epsilon.$$

Then for $z \in \partial G \cap D$,

$$w(z) \geq f(z_0) + \epsilon > f(z),$$

while for $z \in \partial G \setminus D$,

$$w(z) > M \geq f(z).$$

So $w > f$ on the boundary as claimed, and obviously $w(z_0) = f(z_0) + \epsilon$.

It follows that if $v \in \mathcal{P}(f, G)$, then $v \leq w$ on G . Since this is true of all such v , we have $u \leq w$ on G , and so $\limsup_{z \rightarrow z_0} u(z) \leq w(z_0) = f(z_0) + \epsilon$. Since ϵ was arbitrary, $\limsup_{z \rightarrow z_0} u(z) \leq f(z_0)$.

Next let

$$v(z) = f(z_0) - \epsilon - \frac{\phi(z)}{m} [M + f(z_0)] = \left[1 - \frac{\phi(z)}{m} \right] [M + f(z_0)] - M - \epsilon.$$

Now we get $v < f$ on ∂G and $v(z_0) = f(z_0) - \epsilon$. Since $v \in \mathcal{P}(f, G)$, $u \geq v$ on G , and so $\liminf_{z \rightarrow z_0} u(z) \geq f(z_0) - \epsilon$. Letting ϵ tend to zero we have $\liminf_{z \rightarrow z_0} u(z) \geq f(z_0)$. \square

Corollary. *Let G be a bounded region in \mathbb{C} which possesses a barrier at each point of its boundary, and let f be a continuous function on ∂G . Then the Perron function for f on G solves the Dirichlet problem.*

The Schwarz Reflection Principle. To state the reflection principles, we introduce the following terminology: a region G is symmetric with respect to the real axis if $z \in G$ implies $\bar{z} \in G$.

Reflection Principle for Harmonic Functions. *Let G be a region which is symmetric with respect to the real axis and define G_+ , G_- , and G_0 as the intersection of G with the upper half-plane, lower half-plane, and real axis, respectively. If u is a continuous real-valued function on $G_+ \cup G_0$, which is harmonic on G_+ and zero on G_0 , then u admits a unique extension to a harmonic function on all of G . The extension is given by $u(z) = -u(\bar{z})$ for $z \in G_-$.*

Proof. If such an extension exists it is certainly unique, so it suffices to show that the stated extension defines a harmonic function in G . It certainly defines a continuous extension which is harmonic in $G_+ \cup G_-$, so it suffices to show that it is harmonic in a neighborhood of $z_0 \in G_0$. By translating and extending we can assume that $z_0 = 0$ and that G contains the closed unit disk, \bar{D} . Define U on \bar{D} as the solution of the Dirichlet problem with boundary data $u|_{\partial D}$ (where u has been extended to G_- by the formula above). From the Poisson kernel representation we have $U = 0$ on the real interval $(-1, 1)$. Thus $U = u$ on the entire boundary of the upper half-disk, and the entire boundary of the lower half-disk. Since both U and u are harmonic in the half-disks, they coincide, and thus u , like U is harmonic in the whole disk. \square

Reflection Principle for Analytic Functions. *Let G , G_+ , G_- , and G_0 be as above. If f is a continuous complex-valued function on $G_+ \cup G_0$, which is analytic on G_+ and real on G_0 , then f admits a unique extension to a harmonic function on all of G . The extension is given by $f(z) = \overline{f(\bar{z})}$ for $z \in G_-$.*

Proof. We could base a proof on the previous result applied to $\text{Im } f$ and harmonic conjugates, but it is also easy to verify this directly using Morera's theorem. \square

Of course the line of symmetry could be any line, not just the real axis. Thus if G is symmetric with respect to any line, defined and analytic on one side of the line, and real on the line, it can be extended to be analytic on the whole domain. The proof can be found by translating and rotating the domain to the standard case.

We can also translate and rotate the image, so it is not necessary that f be real on the symmetry line, it is sufficient that it map it into some other line. An important application is to analytic continuation. For example, if f maps a rectangle into itself analytically and takes the edges into the edges, we can apply the reflection principle repeatedly to extend f to an entire function.

The reflection principles can be applied for symmetries with respect to circles as well as for lines.

Theorem. *Let f be analytic on the unit disk, continuous on its closure, and real on the boundary. Then f admits a unique extension to the entire plane. The extension is given by $f(z) = \overline{f(1/\bar{z})}$ for $|z| > 1$.*

Proof. Let $g(z) = f((1 + iz)/(1 - iz))$, so g maps the upper half plane into \mathbb{C} and is real on the real axis. Applying the reflection principle to g gives an extension of f with

$$f\left(\frac{1 + iz}{1 - iz}\right) = \overline{f\left(\frac{1 + i\bar{z}}{1 - i\bar{z}}\right)}$$

for z in the lower half plane, or, equivalently,

$$f(w) = \overline{f(1/\bar{w})}$$

for w in the exterior of the disk. \square

As an application of the reflection principle consider an analytic function on the unit disk which extends continuously to the closure taking the unit circle into itself. We may extend this function to \mathbb{C}_∞ by $f(z) = \overline{1/f(1/\bar{z})}$ for $|z| > 1$. (We obtain this by applying the standard reflection principle to $g = \phi^{-1} \circ f \circ \phi$ where $\phi(z) = (1 + iz)/(1 - iz)$.) The extended function is analytic everywhere except at the points symmetric to the zeros of f , where it has a pole. Infinity is either a removable singularity or a pole according to whether f is 0 at 0 or not. Thus the extended function is meromorphic on \mathbb{C}^∞ with a finite number of poles. It follows that f is rational.