

AT MICROFICHE REFERENCE LIBRARY

A project of Volunteers in Asia

Better Farming Series No. 15. Cereals

Published by:

Food and Agriculture Organization of the
United Nations

Via delle Terme di Caracalla

00100 Rome

Italy

Paper copies are \$ 1.50.

Available from:

UNIPUB

P.O. Box 433

Murray Hill Station

New York, NY 10157 USA

Reproduced by permission of the Food and
Agriculture Organization of the United Nations.

Reproduction of this microfiche document in any
form is subject to the same restrictions as those
of the original document.

better farming series

15

1977 edition

cereals



FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS

BETTER FARMING SERIES

Twenty-six titles have been published in this series, designed as handbooks for a two-year intermediate level agricultural education and training course. They may be purchased as a set or as individual documents.

FIRST YEAR

1. The plant: the living plant; the root
2. The plant: the stem; the buds; the leaves
3. The plant: the flower
4. The soil: how the soil is made up
5. The soil: how to conserve the soil
6. The soil: how to improve the soil
7. Crop farming
8. Animal husbandry: feeding and care of animals
9. Animal husbandry: animal diseases; how animals reproduce

SECOND YEAR

10. The farm business survey
11. Cattle breeding
12. Sheep and goat breeding
13. Keeping chickens
14. Farming with animal power
15. Cereals
16. Roots and tubers
17. Groundnuts
18. Bananas
19. Market gardening
20. Upland rice
21. Wet paddy or swamp rice
22. Cocoa
23. Coffee
24. The oil palm
25. The rubber tree
26. The modern farm business

Cereals

**Published by arrangement with the
Institut africain pour le développement économique et social
B.P. 8008, Abidjan, Côte d'Ivoire**

FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS

Rome 1977

First printing 1977

ISBN 92-5-100150-2

©French edition, Institut africain pour
le développement économique et social (INADES) 1970

©English edition, FAO 1977


PREFACE

This manual is a translation and adaptation of "Les céréales", published by the Agri-Service-Afrique of the Institut africain pour le développement économique et social (INADES), and forms part of a series of 26 booklets. Grateful acknowledgement is made to the publishers for making available this text, which it is hoped will find widespread use at the intermediate level of agricultural education and training in English-speaking countries.

The original texts were prepared for an African environment and this is naturally reflected in the English version. However, it is expected that many of the manuals of the series — a list of which will be found on the inside front cover — will also be of value for training in many other parts of the world. Adaptations can be made to the text where necessary owing to different climatic and ecological conditions.

Applications for permission to issue this manual in other languages are welcomed. Such applications should be addressed to: Director, Publications Division, Food and Agriculture Organization of the United Nations, Via delle Terme di Carcalla, 00100 Rome, Italy.

The author of this English version is Mr. A.J. Henderson, former Chief of the Editorial Branch.



OUTLINE OF COURSE

● What are food crops?	2
What cereals are grown in Africa?	3
Why cereals are grown	4
Why we should produce and sell more cereals	5
Traditional cereal growing	6
● Growing cereals in the modern way.	7
Clearing land and grubbing out trees	7
Conserving and improving the soil	9
Applying manure and fertilizers	11
Preparing the soil	12
Selecting and preparing seeds	15
Sowing in rows	17
How to look after cereals	19
Harvesting	23
Selling part of the harvest	27
● Sorghum and millet.	28
Sorghum	28
Millet	30
How to grow sorghum and millet	32
Growing sorghum in the dry season.	35
Sorghum for animal feeding	36
● Maize.	38
Traditional maize growing	40
How to increase maize yields	41
● Other cereals	47
Fonio	47
Finger millet	48
Wheat	49
● Suggested question paper	50

What are food crops?

To live, man must eat.

- At one time men gathered and picked the fruits, leaves, and seeds of trees and plants. To get meat and fish they hunted and fished.
- But nowadays hunting, fishing and food gathering are no longer enough to feed all the people. **Crops have to be grown to provide food.** These crops are called food crops.

- Nowadays the inhabitants of African villages grow many food crops.

They grow mainly:

- **cereals** such as sorghum, millet, rice, maize;
- **tubers** such as yams, sweet potatoes, cocoyams, tania, potatoes;
- **root crops** such as cassava;
- **legumes** such as cow peas, Bambarra groundnuts, groundnuts and soybeans;
- **bananas and plantains.**

*In this booklet we study only cereals.
We shall deal with other food crops
in other booklets.*

What cereals are grown in Africa?

Cereals are plants that yield grains.

Grains are used to feed both man and beast.

- **Certain cereals are grown**
in almost all parts of Africa.
These are mainly millet, sorghum, maize and rice.
Millet and sorghum are grown
in savanna country where it is dry.
Maize is grown everywhere,
in savanna country and in forest country.
Rice used to be grown
only in certain regions.
But now more and more rice is eaten in Africa
and it is grown everywhere.
- **Other cereals are less known.**
They are grown only in some regions.
Fonio is grown
in west Africa.
Finger millet (Eleusine) is grown
in Zaire, Rwanda and Burundi.
Teff is grown
in the mountains of Ethiopia.
Wheat is a cereal of cold countries,
but it grows well in regions near deserts
such as Mauritania, Mali, Niger, Chad,
as well as in hilly regions
like eastern Zaire, Rwanda, Burundi.

*In this booklet we shall deal chiefly with
sorghum, millet and maize.
We shall not speak about rice
because there are two booklets on rice,
one on upland rice and one on wet paddy or swamp rice.
We shall deal briefly with
fonio, finger millet and wheat.*

Why cereals are grown

We grow cereals for their grain.

- **Cereal grains are rich foods.**

They are **energy foods**.

To live, work, walk, men need plenty of energy, plenty of strength. Cereal grains give this strength.

They are **body-building foods**.

Men need to eat such builder foods in order to grow, to make their bodies, their skin and muscles. These builder foods are the proteins. Cereal grains contain lots of protein.

- **Cereal grains are good food**

- **for people.**

The inhabitants of savanna regions live mainly on millet. In all African countries rice has become a very important food.

- **for animals.**

When animal power is used in farm work, the animals have to be well fed.

To make the animals strong, we give them millet or maize as a feed supplement.

Maize can also be given to chickens and pigs.

Cereals can be grown

in order to give their **green stems** to animals.

With the green stems you can make **hay** and **silage** (See the course on animal husbandry).

Why we should produce and sell more cereals

- **We must produce more food to feed more people.**

There are more people in the world.
Fewer children die.
Men and women live longer.

- **We must produce more food to sell to people who do not grow any.**

Many people go to live in towns.
These men and women do not farm the land;
they buy their food.
Even in villages
there are men and women who have new jobs,
such as teachers, officials, traders.
They do not farm the land.
So the farmers have to produce
more millet, more maize, more rice,
to sell to people in towns and villages
who do not farm the land.

- **Farmers should produce more cereals.**

But near the villages
there are often no new fields to cultivate.
In savanna country
there is no time during the short rainy season
to cultivate bigger fields.
So farmers must produce more
from the same area.
**That means they must increase the yield
of their cereal crops.**

**But you cannot increase the yield much
if you farm in the traditional way.**

TRADITIONAL CEREAL GROWING

- In traditional farming
the African farmer clears the land a little,
cuts down the herbage and small trees,
makes a brush fire and sows.
He cultivates this field for two or three years,
then leaves it
and clears another field.
This is called **shifting cultivation**.

This way of farming is bad.

The brush fires spoil the soil
and destroy the organic matter.
Every year the farmer has to clear a new field.
**He loses much time
and often sows too late.**

With shifting cultivation
a great deal of land is unused
and produces no harvest.
Sometimes there is not enough land
to feed the people.
In many places
shifting cultivation cannot be done any more.

- **The farmer works with only a few tools.**
He uses only a machete and a hoe.
He cuts the trees and the herbage with the machete.
He scratches the soil a little with the hoe.
The soil is not well turned over, and water and air
do not get well into the soil.
Plants are not well nourished.
- **The farmer does not enrich the soil.**
Usually he has no animals,
and so does not put any manure on the field.
He does not use fertilizers.
So the soil stays poor and yields a small harvest.

GROWING CEREALS IN THE MODERN WAY

Clearing land and grubbing out trees

A modern farmer who uses animal power
must clear his fields and grub out the trees.

**He must take out all the trees,
even the biggest.**

And he must get out the tree stumps.

If he does not remove the stumps,
his tools (his hoe and plough)
will get bent on the big roots.

He will not be able to use animal power.

He will not be able to sow in rows.

He will not be able to use an animal-drawn cultivator.

The stumps and trees

will go on taking food from the soil.

Around the stumps and trees
the crops will grow less well.

HOW TO GRUB OUT TREES

You can pull out the trees and the stumps
with a tractor.

But that costs a lot,
and most farmers have no tractor.

You can also pull out trees and their stumps
with a winch.

Fix the winch to a big strong tree
and tie the cable of the winch
to the tree you want to pull out.

**Then turn the winch
until the cable pulls out the tree,
stump and all.**

A winch does not cost much.

The farmers in a village can join together
to buy a winch.

Then they can quickly clear
all the fields of the village.


You can also grub out trees with hand tools.

Take away the soil round the base of the tree,
so as to uncover the roots.

Then cut all the roots.

Finally, with a rope pull down the tree.

Its stump will be pulled out at the same time.



Grubbing must be done well before sowing,
at the beginning of the dry season.
At that time the soil is still moist,
and trees are easily pulled out.
Thus the field will be cleared of stumps
before sowing time.
There will be time for tilling the soil
and the sowing will not be late.

Conserving and improving the soil

- **Soil must be protected against erosion**

When a field is cultivated,
the soil is often left bare.

When it rains very hard, the water flows fast,
and carries away the good soil.

This is erosion.

The soil becomes less fertile,
and afterwards the harvests are less good.

When the ground is on a slope,
you must till and make the rows across the slope,
that is, on the **contour lines**.

By this means the water will not flow so fast,
and the good soil will not be carried away.

When the slope is very steep,
terrace farming is used.

You build little walls of earth or stones
to hold back the good soil.

In some mountainous regions
there is a lot of terrace farming.

If the slope is not too steep,
you can make **ditches** along the contour lines,
or **barrier strips**, or **ridges**.

(See the first-year course on the soil).

- **The soil must be protected against the sun.**

When the sun is too strong
it badly damages soil that is left bare.
The sun quickly destroys
the organic material in the soil.
The soil loses its humus
and becomes less fertile.

Bare soil must be covered.

You can sow **cover plants** on land left fallow.
For some crops you can use **mulches** which cover
the soil and enrich it with humus.

- **The fertility of the soil must be conserved.**

To conserve soil fertility, **crop rotation** is used.
**Crop rotation means growing a different crop
each year on the same field.**

Why do we do this?

To feed, plants take
different quantities of mineral salts from the soil.
Different plants have different needs.

They take the mineral salts
at different depths in the soil
because they do not all have the same root system.

With crop rotation,
the plants can use all the mineral salts in the soil
and are better nourished.
The soil does not get so poor.

For example:

after groundnuts grow sorghum.
Groundnuts and sorghum do not have the same
root system, and they have different needs.

*An example of crop rotation is: yams;
cotton;
rice;
groundnuts.*

Applying manure and fertilizers

Manure and fertilizers

**add to the soil
mineral salts which feed the plants,**

**For the plants to make good use of the fertilizers,
you must:**

- **Prepare the soil well.**

**In well-prepared soil, the plant roots develop well;
they can take up all the mineral salts
added by fertilizers and manure.**

- **Sow at the right time.**

**If you sow too late,
the plants do not have enough water
to grow and use all the mineral salts.**

- **Cultivate often.**

**Weeds take up mineral salts.
Weeds must be removed whenever they grow.**

- **Rotate crops.**

**With good crop rotation,
the different plants use all the mineral salts.
The roots take up the mineral salts
at different depths of soil.**

Fertilizers are dear.

**Their cost is not repaid by the harvest,
they are not profitable,
if the farmer does not do all his work well.**

Preparing the soil

Farmers till their soil.

Tilling means turning over the soil.

● Why tilling is needed

Tilling stirs up the soil.

Then water and air can get right into it.

Tilling loosens the soil.

Seeds germinate easily in loose soil,
and roots go well down.

Tilling mixes herbage with the soil.

It cleans the soil.

The herbage rots in the ground and makes humus.

If manure has been spread,
tilling mixes it with the soil.

● The time for tilling

Tilling is best done

at the end of the rainy season,
just after the harvest.

Then the ground will be prepared
and sowing can be done as soon as the rains come.

If tilling is not possible after the harvest,
it must be done as soon as the rains begin,
so that the sowing is not too late.


● **How to till**

Tilling can be done with a **hoe**.

But the work is long and tiring.
The work is not done well;
the soil is not fully turned over.

It is better to till with a **plough**.

You must be a modern farmer,
and till with an animal-drawn plough,
using an ox or donkeys or a horse.
Then the work is done more quickly and better.
The soil is well turned over;
water and air get well into the ground.
You can sow in good time,
you can cultivate bigger fields.



Plough

● Harrowing

After tilling, the soil is often not flat.


There are big pieces of earth, big clods.

To break up the clods, use a **harrow**.


This is a tool with teeth

that break the clods and make the soil flat.

The harrow is drawn by an animal.



If you do not have a harrow,
make an animal drag **large branches** over the soil.
The branches break up and crush the clods.



Selecting and preparing seeds

To get a good harvest,
you must sow good seeds.

- **Farmers can buy selected seeds.**

Research stations select the best varieties,
the ones which are best adapted to the climate,
which resist diseases well
and give plentiful harvests.

These seeds are dear.
To pay for them, you need a good harvest.
The harvest will be good
only if you do all your work well.

- **Farmers can select seeds from their own crops.**

Take the finest heads of grain,
and from them only the well-formed seeds.
The good qualities of these seeds
will be passed on to the plants that grow from them.

- **Farmers must store their seeds well.**


Seeds should be dried
before being put into the granary.
Seeds which are not quite dry may rot.

Cereal seeds must be protected
against insects and rats.
To protect seeds against insects,
the seeds are mixed with an insecticide.
To protect seeds against rats,
the granary is raised well above the ground.

● Preparing seeds

Before sowing,
take away all seeds that are broken,
bitten, diseased or misshapen.

The seeds may also be **disinfected**.
Mix the seeds with a pesticide,
and see that the seeds are well covered by it.



Device for treating seeds

Disinfected seeds are not eaten by insects.
They do not rot.
All the seeds will grow
and very few plants will be missing in the rows.

**By sowing disinfected seeds
you get a good density;
the yield is better.**


Sowing in rows

In Africa it is usual to sow cereals broadcast.
This is a bad way of doing it.

- **It is better to sow in rows.**

Sowing in rows means that the seedlings will be at the same distance from each other. Animal power can be used and the cultivations are easier; the animal that pulls the hoe can pass between the rows. It is easier to give each plant the same amount of fertilizers.

On flat soil the rows can be traced with a marker.
The marker can be pulled by hand or by a donkey.
The spikes of the marker trace the rows where the seed will be planted.




When the seed rows
have been traced by the marker,
the farmer puts the cereal seeds
the same distance apart along each row.

- To do the work more quickly,
a **seed drill** can be used.
It is pulled by an animal.
The seed drill makes a furrow in the soil
and drops the seeds
at the same distance from each other.
A little wheel at the back of the drill
covers the seeds with 2 or 3 centimetres of earth.

With this machine you can **sow quickly**
and with an **even density**.

Some seed drills apply fertilizer
at the same time.



Seed drill

HOW TO LOOK AFTER CEREALS

- **Seedlings which have not grown must be replaced.**

A week after sowing you will see empty places where no plants have grown. Sow again in these empty places, so as to have a fully planted field of good density.

- **Thinning must be done.**

If you have sown in seed holes, there are often 4 or 5 young plants in each seed hole. To make the cereals grow better, it is best not to keep all the seedlings. Keep only two seedlings in each hole. Pull out the smallest, and keep the best plants. When you pull out the unwanted seedlings, you disturb the soil. So you must press the soil back round the young plants that remain.

With one hand, hold the soil round the plants that are not removed.



CULTIVATING

Cultivating means removing weeds .

- **Why cultivating is needed**

Weeds in the ground take food away from the crop.
They take water and mineral salts
out of the soil.

They also take the mineral salts
added in fertilizers.

Weeds take the place of good plants.
They cast shade.

By cultivating, you stir up the soil.
You mix air with the soil,
break up the dry crust,
and prevent water from rising and being lost.
The soil remains moister.

- **When to cultivate**

The first cultivation
must be done when the weeds are still small,
two weeks after sowing.
This first cultivation is very important,
because it enables the young plants to make a good start.

Afterwards you must cultivate twice more,
each time when the weeds have grown.

When you cultivate you can also do the **earthing up** .
At the base of the cereal stem
there are adventitious roots.
To make these roots grow well,
earth must be placed around the foot of the stem.
This is called earthing up.
Do the earthing up
at the same time as the first cultivation.

● **How to cultivate**

You can cultivate
with a **hand hoe**
or with an **animal-drawn cultivator**.

If the cereals are sown in rows,
you can work **more quickly**
by using an animal-drawn cultivator.

The work is well done

and you can cultivate **more often**.

The cultivator uproots the weeds between the rows.

Afterwards you must go with the hand hoe

to take out the weeds in the rows.


Leave uprooted weeds on the ground.

They protect the soil

against erosion

and against the sun.

They rot and form humus.



Animal-drawn cultivator

CONTROLLING THE ENEMIES OF CEREAL CROPS

The chief enemies of cereals
are **animals**,
and above all, **birds**.

- **Goats, sheep and cows**

go into fields of millet, maize and rice
and eat the young plants.

You can surround the fields with a fence
so that the animals cannot get in.

But the best way is to **shut up all the animals**.

Put the oxen and cows in one paddock,
and the sheep and goats in another.

Then your fields will not be spoiled.

- **But the most dangerous enemies are birds.**

First of all, they may eat the seeds in the earth,
just after sowing.

But above all, they come and eat the grains in the ear,
when the crop is ripe.

Usually children guard the fields
and chase the birds away.

But nowadays children go to school.

You can put **scarecrows**,
dressed to look like a man,
in the fields.

The birds are frightened
and do not come and eat the grain.

- Different **insects** may also attack cereals, such as
caterpillars,
grasshoppers,
aphids.

You can control insects
by using a pesticide such as BHC.

HARVESTING

- **Cutting the stems of cereals**

When the grain is ripe,
you must not wait too long before harvesting.
If you wait too long,
the grain may fall to the ground,
and birds may come and eat the grain on the plant.

Cut the stems
with a machete, a sickle or a knife.



- Afterwards cut off the heads of grain with a knife.
Leave the stems on the ground.
These stems cover the ground
and protect the soil from the sun and from erosion.
After rotting,
the stems add organic matter to the soil.
If you keep animals,
you can put the stems in their stalls or paddock.
Mixed with the animal droppings,
the stems make manure which will enrich the fields.

- **Drying the grain**

When you have cut the heads,
you must not thresh them at once,
nor put them in a granary.
First of all, you must dry them very well.
Spread the heads on a really clean surface.
Stop sheep, goats and birds
from eating the grain.


When the heads are well dried,
you can thresh them,
that is, separate the grain from the heads.
But do not thresh all your crop at once.
Keep the greater part of the harvest
as heads of grain.
Thresh only the amount you need for food.

- **Threshing cereals**

**Threshing means
getting the grain out of the heads.**

Threshing can be done
by beating the heads with a stick,
or with a special tool called a **flail**.
For rice there is a machine
that separates the rice grains from the heads.
It is called a **thresher**.

**Small rice
thresher**



- **Winnowing the grain**

Cereal grain must be very clean.

This is important.

It must not be mixed
with earth and little stones.

When the grain has been threshed,
it is winnowed to make it cleaner.

For winnowing, use a sieve,
or else pour the grain
from one flat bowl into another.



- **Storing cereal grain**


Storing can be done in sacks or in granaries.

The sacks and the granaries must be protected against damp, which makes the grain rot, and against rats and insects, which eat or spoil the grain.

The granaries must be above the ground to keep the grain dry.

The granaries must be disinfected.

Ask the extension officer what pesticides to use and how to apply them. Some pesticides are poisonous.



Clay granary in northern Cameroon

SELLING PART OF THE HARVEST

- Often farmers sell millet or maize just after the harvest.
At that time, there is plenty of millet in all the villages.
Nobody needs to buy any.
The price of millet is very low.
Farmers who sell their grain just after the harvest do not earn much money.
- Sometimes, at the beginning of the rainy season, these farmers have no millet left for food.
At that time, there is hardly any millet in the granaries.
Millet is very dear.
Farmers buy a little millet, while waiting for the harvest.
But they pay a high price.
- It is better not to sell your millet or maize at harvest time,
for the selling price is low,
and you will perhaps not have enough grain for food all through the year.
- Keep your harvest safe in your granaries.
You can then sell your grain at the beginning of the rainy season, when food is beginning to get short.
Then you will earn more money, for the price of millet or maize is much higher.

Sell when the price is high .

SORGHUM AND MILLET

Sorghum and millet are very important cereals in savanna country.

They are grown in the same way.

We shall deal with the growing of sorghum and millet together.

SORGHUM




Sorghum plant

Grain

Sorghum is a tall cereal, more than 2 metres high.

Its stem has long leaves.

At the top of the stem is the spike.



Spike of sorghum

The spike contains the sorghum grains. Their colour is white, black, yellow or red.

Sorghum grows in savanna country that is not too dry.

It needs about 1 metre of rain a year.

Sorghum likes light soil that lets rain run through, that is, permeable soil.

VARIETIES OF SORGHUM

Sorghum is sometimes called "great millet" because it has rather large grains.

- There are several very different varieties of sorghum:
 - **Grain sorghums**
These are the kind most grown in Africa.
They yield grain for human food.
 - **Sweet sorghums**
The stem contains a sweet liquid.
They are grown in equatorial Africa
and in northern Cameroon.
 - **Broom sorghums**
These have very large spikes, like a broom.
 - **Fodder sorghums**
These are cut when still green
for animal feeding.
- Each of these varieties includes several kinds that ripen at different times.
 - **Early sorghums** ripen 80 to 110 days after sowing.
 - **Medium early sorghums** ripen 115 to 130 days after sowing.
 - **Medium late sorghums** ripen 130 to 145 days after sowing.
 - **Late sorghums** ripen about 190 days after sowing.

Some kinds of sorghum are preferred for **human food**, others for making **beer**.

MILLET



Millet plant

- Millet is a tall cereal, more than 2 metres high.

Its stem has long leaves. At the top of the stem is the spike.

It is not like the spike of sorghum.

The spike contains the millet grains. They are smaller than sorghum grains. They are white or red in colour.

- Millet grows in very dry regions north of the savanna. It even grows where the rainfall is only **300 millimetres a year**.

Millet likes light, sandy and permeable soils.

VARIETIES OF MILLET

Different names for millet are
bulrush millet, pearl millet,
Japanese millet, broomcorn millet.

- There are a great many varieties of millet.
They have different names in different countries,
for instance:
 - in Senegal: *Souna, Sanio*;
 - in northern Cameroon: *Youri, Yadiri*;
 - in Chad: *Dokone*.


- The different kinds of millet
are distinguished by the length of the spike.
 - Varieties with a long spike
In Niger there is *Rongo*,
with spikes that may be more than 2 metres long,
and *Somno*,
with spikes 80 centimetres long.
These varieties are usually late ones
that ripen 120 to 150 days after sowing.

 - Varieties with medium or cylindrical spikes
The length of the spike is about 40 centimetres.
They are harvested about 100 days after sowing.

 - Varieties with short spikes
For instance, in Niger,
Batoutchani has spikes of 6 centimetres,
Boudouma, spikes of 10 to 20 centimetres.
These are usually early varieties.
They ripen 60 to 90 days after sowing.

How to grow sorghum and millet

- Sorghum and millet have fibrous roots.



To be well nourished
and resist dryness,
sorghum and millet need
well-developed roots.

The roots of sorghum and millet are thin.
They are not strong enough
to penetrate hard earth,
soil that is much compacted, and clods.
They must have **fine, well-loosened soil,**
without clods.

After tilling,
break up the clods with a harrow
or large branches drawn by an animal.

If the soil is not deep,
make little mounds or ridges.

● The place of sorghum in crop rotation

It is best to sow sorghum
after growing another crop
such as groundnuts or cotton.

If sorghum is grown after groundnuts or cotton,
it will use the remainder of the mineral salts
added for these industrial crops.

HOW TO SOW

Selected your seed.

You can keep the seed
from the best spikes of your own harvest.
Dry them, sort them
and store them well,
protect them against insects and diseases.

You can also buy seed of improved varieties
from research stations.

Some varieties are harvested
about 100 days after sowing.
These varieties are sown
in regions where the rainy season is short.

Other varieties are harvested
150 to 190 days after sowing.
These varieties are sown
in regions where the rainy season is longer.

Sow at the beginning of the rainy season.

Sow in rows,
and leave 80 to 100 centimetres between rows.
Put 4 to 10 seeds in each seed hole,
and leave 40 to 60 centimetres between seed holes.
Push the seed 2 centimetres into the soil.

CARE OF THE CROP

Weed the field
two weeks after sowing.

In seed holes where the seeds have not grown,
new seed can be sown,
or seedlings can be transplanted.
Take them from seed holes
where all the seeds have grown.

In seed holes where all the seeds have grown,
take out the smallest seedlings.
It is better to have fewer but finer plants
with bigger spikes.
Then there will be more grain.

When the plants are 50 centimetres high,
weed again.

At the same time,
earth up the plants.
After earthing up, new roots will grow.

HARVESTING

When the grain is ripe,
do not wait too long
before harvesting sorghum and millet.
If you wait too long,
birds may eat a lot of the grain.

After harvesting,
you can thresh the spikes
and store the sorghum and millet grain.
But you can also store the spikes
with the grain in them.

GROWING SORGHUM IN THE DRY SEASON

- In countries such as Niger, Cameroon and Chad, sorghum is sown along rivers and streams and by the side of lakes at the end of the rainy season.

When the rain stops, the water subsides along the river banks and lake sides, but the soil, which is rather clayey, retains water and remains **moist**. Sorghum is then sown on this moist earth. The sorghum grows during the dry season by taking up the water that remains in the soil. The sorghum becomes ripe during or at the end of the dry season. Thus, there is a sorghum harvest just before the **hungry season**.

Sorghum can also be started in **nurseries**. The seed is sown on a little plot which is watered. At the end of the rainy season when the water subsides, the young sorghum plants are lifted and transplanted to river banks or lake sides. The sorghum ripens in the middle of the dry season.

Sorghum grown during the dry season is called *Berbéré* in Chad and *Mouskouari* in northern Cameroon.

SORGHUM FOR ANIMAL FEEDING

When young sorghum plants are still green,
they may be fed to animals.

But you must be very careful,
because sorghum may contain a poison.
The quantity of poison in the plants
is not always the same.
It differs according to the age of the plants,
the variety, the soil, the climate.

Sometimes when you have cut the green sorghum,
some plants may produce new stems.
These new growths contain a lot of poison.

**So take good care
before giving green sorghum to animals.**

The animals can be fed sorghum
by putting them in the field where it is growing,
or by giving it to them
as hay
or as silage.

Never give the new growths to animals.

- **Feeding animals in the fields**

Animals are let into the field
when the spikes of sorghum begin to ripen.

- **Making sorghum hay**

To make hay,
cut the sorghum plants
when the spikes have come out of the stem.
Leave the stems and leaves to dry.

**Because of the poison,
you must wait two months
before giving this hay to animals.**

- **Making silage**

To make silage,
cut the sorghum plants
when the spikes are well formed,
but before the grains have become hard.
Then wait two months
before giving the silage to animals.

**Never make silage
with new growths of sorghum.**

There are some varieties of sorghum
that are grown only as fodder.
These varieties are very good for making silage.

MAIZE


VARIETIES OF MAIZE

- **There are many varieties of maize:**
 - **Soft maize**
The grain has no hard husk,
and it is very floury;
 - **Hard maize**
The grain has a hard husk.
It can be kept for a fairly long time.
 - **Sweet maize**
It is eaten before ripening.
- **There is also the difference in colour
between white maize and yellow maize.**

**When the maize plants are still green,
they can be used as fodder.**
Silage can also be made from them.

Some parts of Africa could produce more maize
to feed the growing population
and to feed the animals better.

Maize could also be grown
to sell to foreign countries.



Male flowers

Female flowers

Maize plant

In a maize plant,
stamens and ovaries
are in different
flowers.

TRADITIONAL MAIZE GROWING

- Traditionally, maize is grown in association with other crops.

It is grown in the same field with yams, cotton or groundnuts. When several crops are grown in the same field, harvests are poor, sowing is not done at the right time, weeding is not well done, and much time is lost in harvesting.

- Traditionally, maize is grown without the use of new tools.

The soil cannot be well worked with the hand hoe, and so the soil is not loosened; the roots cannot penetrate well into the soil to take up water and mineral salts; the plants are not well nourished. There is not much grain; the yield is low.

- Traditionally, maize is grown without fertilizers.

No mineral salts are added to the soil; the soil becomes poor. So old fields must be left fallow and new fields have to be cleared. This is **shifting cultivation**.

With shifting cultivation, much land is not cultivated and yields no harvest. Farmers lose a lot of time clearing new fields.

Shifting cultivation must stop in order to increase maize production.


HOW TO INCREASE MAIZE YIELDS

● Choosing the seeds

If you want a good harvest,
you must sow good seeds.

You can choose seeds from your own crop.
When you harvest, choose the biggest cobs.
Keep these cobs.
For sowing,
**take only the best-shaped grains
from the centre of the cobs.**
If you choose your seeds in this way,
you will have a better harvest.
Seed dressing is also useful.

The finest grains from
the centre of the cob
are chosen



But yields become smaller
if you sow every year
seeds from last year's crop.
After a few years,
the maize loses its good qualities.
So you must buy new seeds.

- The research stations have improved varieties of maize which produce very good seeds.


Several varieties of maize are grown at the research stations. The female flowers of one variety are fertilized with the pollen from the male flowers of another variety to produce hybrids. Otherwise, selection breeding is carried out within the same variety.



Pollen from the male flowers of one variety

fertilizes

the female flowers of another variety



The little bag prevents pollen from fertilizing the female flowers of the same plant

When the female flowers of one maize variety are fertilized by pollen from the male flowers of another maize variety, this is called **crossing**. The flowers fertilized in this way produce a new variety of maize called a **hybrid**, which is better than the two original varieties. But after several years of growing maize from the previous crop's seeds, this new variety loses its good qualities. So new seeds have to be bought.

Crossing two varieties may also result in new varieties that are more resistant to disease.

- **Preparing the soil**

Maize needs fine earth,
without clods, a loose soil,
in order to develop its root system.
Tilling with a plough loosens the soil.
The clods are broken up with the harrow.

HOW TO SOW

Sow your maize at the beginning of the rainy season.

Sow in rows,
leaving 80 centimetres between rows.
Put 3 to 4 seeds in one hole,
leaving 50 centimetres between holes.

Make each hole 7 centimetres deep
and cover the seeds with a little earth.
Like that, the seeds have more protection
against rats and birds.

HOW TO LOOK AFTER MAIZE

- **Weeding is very important.**

This work can be done better and more quickly
with an animal-drawn cultivator.

Weed for the first time
15 days after sowing.

Pull out the smallest seedlings
from holes where all the seeds have come up;
leave only the largest seedling.

Weed for the second time
when the plants are 50 centimetres high.
When you weed, also earth up the plants.
After earthing up,
the plants will have more adventitious roots,
and the maize will be more resistant to drought.
The maize will be better nourished.

- **Manure and chemical fertilizers**

Organic manure and chemical fertilizers cannot be used well and it is not profitable to apply them:

- if the soil is not well prepared,
- if the seeds are not well chosen,
- if sowing is not done at the right time,
in rows and at a good density,
- if the weeds are not removed whenever they grow,
- if the crop rotation is not good.

- **Different manures and fertilizers have to be applied:**

- **according to your crop rotation.**

Different plants do not all take the same amount of each mineral salt out of the soil. Before sowing maize, you must therefore give back to the soil the mineral salts taken out by the previous crop. These mineral salts are contained in the fertilizers.

- **according to the region.**

The soils of different regions do not all contain the same amount of each mineral salt. They must therefore be given different fertilizers.

For example:

If a soil is poor in nitrogen, the tips of the young leaves turn yellow. You must therefore add nitrogen to the soil.

You can use:

- **organic manures**

These are animal manure and green manure.

Organic manure improves the structure of the soil. Plants grow better in a soil of good structure, and the chemical fertilizers are used better. Organic manuring should therefore be done at the beginning of the rotation, that is, before growing the first crop on a field.

- **chemical fertilizers**

Maize needs above all nitrogen.

The nitrogen fertilizer most often used in Africa is ammonium sulphate.

But maize also needs phosphate and potassium.

Ask the extension services for advice.

They will tell you how much fertilizer to apply to your maize field.

For example:

At Bouaké (Ivory Coast), the following fertilizers are applied before sowing one hectare of maize:

100 kg ammonium sulphate,

100 kg dicalcium phosphate,

50 kg potassium chloride

After the first cultivation, another 100 kg ammonium sulphate may be added. Be careful not to put any fertilizer on the leaves; the fertilizer may burn the leaves.

HARVESTING

Maize can be fed to animals
when the plants are still green.
In that case, cut the maize
when the male flowers begin to go dry.
You can also make silage.

If you want to roast the maize cobs,
harvest the cobs
when the grains are not too ripe.

If you want to make flour,
harvest the cobs
when the grains are quite ripe.

STORING MAIZE

Maize cobs can be stored in their husks.
That way, the cobs keep better.

They can also be stored without their husks,
but then the cobs
may be attacked by insects.
There are insecticides
which kill these insects.



Maize cob in husk



Maize cob without husk

OTHER CEREALS

FONIO

Fonio is a **short cereal**,
about 45 centimetres high.
It yields very small grains.

Fonio is chiefly grown
in **savanna regions** of west Africa,
such as Senegal, Mali, Guinea, Upper Volta, Niger.

Fonio grows and ripens quickly.
It can be harvested long before millet or sorghum.
Thus, the inhabitants of savanna regions
who have no millet or sorghum during the hungry season
can eat fonio.

Fonio grows well even on **poor soil**.
It is much grown in certain mountainous regions
such as Fouta Djallon in Guinea,
and on soils with a little laterite
in Upper Volta and Mali.
It is a plant that needs very little water to grow well.
It is grown on very dry and poor soils.

Fonio should be sown at the end of a rotation,
just before the fallow.

Fonio fields must be protected
from **birds** that eat the grain.

Harvesting fonio is difficult,
because the grain is very small.
Much grain can be lost in harvesting.
Do not wait until the grain is too ripe.

FINGER MILLET

Finger millet (*Eleusine*)

grows to between 60 and 150 centimetres high.

It is grown in Zaire, in the Central African Empire, in Rwanda, in Burundi, in Ethiopia, in Kenya, and in Tanzania.

At the top of the stem

there are **5 to 10 ears**;

they are curved and contain the grain.

Finger millet needs **plenty of water**.

It is chiefly grown in **hilly regions**,
on **high tablelands**.

Finger millet grows well

even on poor soil.

It is often grown together with other crops
such as maize and groundnuts,
or after other crops such as cotton.

The ears are harvested by hand

and put in bundles to dry.

Then they are stored in granaries.

The ears store well,
and are not attacked by insects.

Finger millet is used for food;

flour made from it is cooked.

It is also eaten in the form of **paste**.

Some kinds of finger millet are used for making **beer**.

WHEAT

Wheat is a cereal

that is chiefly grown in **cold countries**.

Flour is made from wheat
and is used for making **bread**.

In Africa, especially in towns,
people eat more and more bread.

Until now, African countries
bought wheat or wheat flour from foreign countries.

But that is costly.

It is better, when possible,
to produce the wheat needed for food.

In Africa, wheat is grown

**in regions where there is a very cold season,
that is:**

- **on the edge of the desert,
near rivers and lakes:**
 - in Mauritania and Senegal,
along the Senegal river;
 - in Mali and Niger,
along the Niger river;
 - in Chad,
near Lake Chad;
- **in mountainous and hilly regions
such as:**
 - eastern Zaire;
 - Rwanda and Burundi.

**Some countries, such as Chad,
are making a great effort to develop wheat growing.
This is being done chiefly in the Lake Chad region
which will soon be able to meet
all the country's needs of wheat.
Chad will no longer have to import wheat.**

SUGGESTED QUESTION PAPER

FILL IN THE MISSING WORDS

Cereals are plants which yield

When the slope is very steep is used.

Crop rotation means growing a each year on

Farmers who sell their grain just after the harvest money.

Cereal grains are good food for and for

Millet stems are cut with a

Grain should be sold when

Wheat flour is used for making

ANSWER THE FOLLOWING QUESTIONS

What cereals are grown in your region?

Where you live are cereal grains given to animals?

Why must more cereals be produced?

What is shifting cultivation?

Why must trees be grubbed out?

Where you live is a winch used for grubbing?

How do you protect the soil against the sun?

What is thinning?

Which are the chief enemies of cereals?

How do they thresh cereals where you live?

When should the farmer sell his millet? Why?

What are the different names of millet in the language of your country?

What is each variety of millet used for (food, beer, ...)?

What is the way to get a new variety of maize?

Why must you sow at the right time?

Are fonio, finger millet or wheat grown where you live?

FAO SALES AGENTS AND BOOKSELLERS

Antilles, Netherlands	Boekhandel St. Augustinus, Abraham de Veerstraat 12, Willemstad, Curaçao.
Argentina	Editorial Hemisferio Sur S.R.L., Libreria Agropecuaria, Pasteur 743, Buenos Aires.
Australia	Hunter Publications, 58A Gipps Street, Collingwood, Vic. 3066; The Assistant Director, Sales and Distribution, Australian Government Publishing Service, P.O. Box 84, Canberra, A.C.T. 2600, and Australian Government Publications and Inquiry Centres in Canberra, Melbourne, Sydney, Perth, Adelaide and Hobart
Austria	Gerold & Co., Buchhandlung und Verlag, Graben 31, 1011 Vienna.
Bangladesh	Agricultural Development Agencies in Bangladesh, P.O. Box 5045, Dacca 5.
Barbados	Cloister Bookstore Ltd., Hincks Street, Bridgetown.
Belgium	Service des publications de la FAO, M.J. De Lannoy, rue du Trône 112, 1050 Brussels. CCP 000-0808993-13.
Bolivia	Los Amigos del Libro, Perú 3712, Casilla 450, Cochabamba; Mercado 1315, La Paz; Rene Moreno 26, Santa Cruz; Junin esq. 6 de Octubre, Oruro.
Brazil	Livraria Mestre Jou, Rua Guaipa 518, São Paulo 10; Rua Senador Dantas 19-S205/206, Rio de Janeiro.
Brunei	MPH Distributors Sdn. Bhd., 71/77 Stamford Road, Singapore 6 (Singapore).
Canada	Renouf Publishing Co. Ltd., 2182 Catherine St. West, Montreal, Que H3H 1M7.
Chile	Biblioteca, FAO Oficina Regional para América Latina, Av. Providencia 871, Casilla 10095, Santiago.
China	China National Publications Import Corporation, P.O. Box 88, Peking.
Colombia	Litexsa Colombiana Ltda., Calle 55, N° 16-44, Apartado Aéreo 51340, Bogotá.
Costa Rica	Libreria, Imprenta y Litografía Lehmann S.A., Apartado 10011, San José.
Cuba	Instituto del Libro, Calle 19 y 10, N° 1002, Vedado.
Cyprus	MAM, P.O. Box 1722, Nicosia.
Denmark	Ejnar Munksgaard, Norregade 6, Copenhagen S.
Dominican Rep.	Fundación Dominicana de Desarrollo, Casa de las Gárgolas, Mercedes 4, Santo Domingo.
Ecuador	Su Libreria Cia. Ltda., Garcia Moreno 1172, Apartado 2556, Quito.
El Salvador	Libreria Cultural Salvadoreña S.A., Avenida Morazán 113, Apartado Postal 2296, San Salvador.
Finland	Akateeminen Kirjakauppa, 1 Keskuskatu, Helsinki.
France	Editions A. Pedone, 13 rue Soufflot, 75005 Paris.
Germany, F.R.	Alexander Horn Internationale Buchhandlung, Spiegelgasse 9, Postfach 3340, Wiesbaden.
Ghana	Fides Enterprises, P.O. Box 1628, Accra.
Greece	« Eiefttheroudakis », 4 Nikis Street, Athens.
Guatemala	Distribuciones Culturales y Técnicas « Artemis », Quinta Avenida 12-11, Zona 1, Guatemala City.
Guyana	Guyana National Trading Corporation Ltd., 45-47 Water Street, Georgetown.
Haiti	Max Bouchereau, Librairie « A la Caravelle », B.P. 111B, Port-au-Prince.
Honduras	Editorial Nuevo Continente S. de R.L., Avenida Cervantes 1230-A, Apartado Postal 380, Tegucigalpa.
Hong Kong	Swindon Book Co., 13-15 Lock Road, Kowloon.
Iceland	Snaebjörn Jónsson and Co. h.f., Hafnarstraeti 9, P.O. Box 1131, Reykjavik.
India	Oxford Book and Stationery Co., Scindia House, New Delhi; 17 Park Street, Calcutta.
Indonesia	P.T. Gunung Agung, 6 Kwitang, Djakarta.
Iran	Iran Book Co. Ltd., 127 Nadershah Avenue, P.O. Box 14-1532, Tehran; Economist Tehran, 99 Sevom Estand Avenue, Tehran (sub-agent).
Iraq	National House for Publishing, Distributing and Advertising, Rashid Street, Baghdad.
Ireland	The Controller, Stationery Office, Dublin.
Israel	Emanuel Brown, P.O. Box 4101, 35 Allenby Road and Nachlat Benyamin Street, Tel Aviv; 9 Shlomzion Hamalka Street, Jerusalem.
Italy	Distribution and Sales Section, Food and Agriculture Organization of the United Nations, Via delle Terme di Caracalla, 00100 Rome; Libreria Scientifica Dott. L. De Blasio « Aeiou », Via Meravigli 16, 20123 Milan; Libreria Commissionaria Sansoni « Licoso », Via Lamarmora 45, C.P. 552, 50121 Florence.
Jamaica	Teachers Book Centre Ltd., 96 Church Street, Kingston.
Japan	Maruzen Company Ltd., P.O. Box 5050, Tokyo Central 100-31.
Kenya	Text Book Centre Ltd., P.O. Box 47540, Nairobi.
Korea, Rep. of	The Eul-Yoo Publishing Co. Ltd., 5 2-Ka, Chong-ro, Seoul.

FAO SALES AGENTS AND BOOKSELLERS

Kuwait	Saeed & Samir Bookstore Co. Ltd., P.O. Box 5445, Kuwait.
Lebanon	Dar Al-Maaref Liban S.A.L., place Riad El-Solh, B.P. 2320, Beirut
Luxembourg	Service des publications de la FAO, M.J. De Lannoy, rue du Trône 112, 1050 Brussels (Belgium).
Malaysia	MPH Distributors Sdn. Bhd., 9A Jalan 14/20, Section 14, Petaling, Jaya.
Mauritius	Nalanda Company Limited, 30 Bourbon Street, Port Louis.
Mexico	Dilitsa, Puebla 182-D, Apartado 24-448, Mexico City 7, D.F.
Morocco	Librairie « Aux Belles Images », 281 avenue Mohammed V, Rabat
Netherlands	N.V. Martinus Nijhoff, Lange Voorhout 9, The Hague
New Zealand	Government Printing Office, Government Bookshops at Rutland Street, P.O. Box 5344, Auckland; Mulgrave Street, Private Bag, Wellington, 130 Oxford Terrace, P.O. Box 1721, Christchurch, Princes Street, P.O. Box 1104, Dunedin, Alma Street, P.O. Box 857, Hamilton.
Nicaragua	Incusa-Culturama, Camino de Oriente, Apartado C105, Managua
Nigeria	University Bookshop (Nigeria) Ltd, University of Ibadan, Ibadan
Norway	Johan Grundt Tanum Bokhandel, Karl Johansgt. GT 41-43, Oslo 1.
Pakistan	Mirza Book Agency, 65 The Mall, Lahore 3
Panama	Distribuidora Lewis S.A., Edificio Dorasol, Calle 25 y Avenida Balboa, Apartado 1634, Panama 1.
Peru	Libreria Distribuidora Santa Rosa, Jirón Apurímac 375, Lima
Philippines	The Modern Book Company, 928 Rizal Avenue, Manila.
Poland	Ars Polona-Ruch, Krakowskie Przedmiescie 7, Warsaw
Portugal	Livraria Bertrand, S.A.R.L., Apartado 37, Amadora; Livraria Portugal, Dias y Andrade Ltda., Apartado 2681, Rua do Carmo, 70-74, Lisbon-2; Edicoes ITAU, Avda. República 46A c/v-E, Lisbon-1
Romania	Hlexim, Calea Grivitei, N° 64-66, B.P. 2001, Bucarest.
Saudi Arabia	University Bookshop, Airport Road, P.O. Box 394, Riyadh
Senegal	Librairie Africa, 58 Av. Georges Pompidou, B.P. 1240, Dakar
Singapore	MPH Distributors Sdn Bhd., 71/77 Stamford Road, Singapore 6
Somalia	« Samater's », P.O. Box 936, Mogadishu.
Spain	Mundi Prensa Libros S.A., Castelló 37, Madrid 1; Libreria Agricola, Fernando VI 2, Madrid 4.
Sri Lanka	M.D. Gunasena and Co. Ltd., 217 Norris Road, Colombo 11.
Switzerland	Librairie Payot S.A., Lausanne et Genève; Buchhandlung und Antiquariat, Heinimann & Co., Kirchgasse 17, 8001 Zurich.
Surinam	VACO nv in Surinam, P.O. Box 1841, Domineestraat 26/32, Paramaribo
Sweden	C.E. Fritzes Kungl. Hovbokhandel, Fredsgatan 2, 103 27 Stockholm 16
Tanzania	Dar us Salaam Bookshop, P.O. Box 9030, Dar es Salaam.
Thailand	Suksapan Panit, Mansion 9, Rajadamnern Avenue, Bangkok
Togo	Librairie du Bon Pasteur, B.P. 1164, Lomé
Trinidad and Tobago	The Book Shop, 111 Frederik Street, Port of Spain.
Turkey	Güven Bookstores, Güven Bldg., P.O. Box 145, Müdafaa Cad. 12/5, Kizilay-Ankara; Güven Ari Bookstores, Ankara Cad. No. 45, Cagaloglu-Istanbul; Güven Bookstore, S.S.K. Konak Tesisleri P-18, Konak-Izmir.
United Kingdom	Her Majesty's Stationery Office, 49 High Holborn, London WC1V 6HB (callers only); P.O. Box 569, London SE1 9NH (trade and London area mail orders); 13a Castle Street, Edinburgh EH2 3AR; 41 The Hayes, Cardiff CF1 1JW; 80 Chichester Street, Belfast BT1 4JY; Brazennose Street, Manchester M60 8AS; 258 Broad Street, Birmingham B1 2HE; Southey House, Wine Street, Bristol BS1 2BQ.
United States of America	UNIPUB, 345 Park Avenue South, New York, N.Y. 10010; mailing address: P.O. Box 433, Murray Hill Station, New York N.Y. 10016.
Uruguay	Juan Angel Peri, Alzarbar 1328, Casilla de Correos 1755, Montevideo.
Venezuela	Blume Distribuidora S.A., Av. Rómulo Gallegos esq. 2a. Avenida, Centro Residencial « Los Almendros », Torre 3, Mezzanina, Ofic. 6, Urbanización Montecristo, Caracas.
Yugoslavia	Jugoslovenska Knjiga, Terazije 27/11, Belgrade; Cankarjeva Založba, P.O. Box 201-IV, Ljubljana; Prosveta Terazije 16, P.O. Box 555, 11001 Belgrade.
Other countries	Requests from countries where sales agents have not yet been appointed may be sent to: Distribution and Sales Section, Food and Agriculture Organization of the United Nations, Via delle Terme di Caracalla, 00100 Rome, Italy.