# INTERNATIONAL STANDARD

**ISO** 9842

Second edition 2003-09-01

# Oil of rose ( $Rosa \times damascena$ Miller)

Huile essentielle de rose (Rosa x damascena Miller)





#### PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

#### © ISO 2003

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office Case postale 56 • CH-1211 Geneva 20 Tel. + 41 22 749 01 11 Fax + 41 22 749 09 47 E-mail copyright@iso.org Web www.iso.org

Published in Switzerland

#### **Foreword**

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 9842 was prepared by Technical Committee ISO/TC 54, Essential oils.

This second edition cancels and replaces the first edition (ISO 9842:1991), which has been technically revised.

### Oil of rose (Rosa × damascena Miller)

#### 1 Scope

This International Standard specifies certain characteristics of the oil of rose ( $Rosa \times damascena$  Miller) cultivated in Turkey, Morocco and Bulgaria, of the Rosaceae family, in order to facilitate assessment of its quality.

#### 2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO/TR 210, Essential oils — General rules for packaging, conditioning and storage

ISO/TR 211, Essential oils — General rules for labelling and marking of containers

ISO 212, Essential oils — Sampling

ISO 279, Essential oils — Determination of relative density at 20 °C — Reference method

ISO 280, Essential oils — Determination of refractive index

ISO 592, Essential oils — Determination of optical rotation

ISO 709, Essential oils — Determination of ester value

ISO 1041, Essential oils — Determination of freezing point

ISO 11024-1, Essential oils — General guidance on chromatographic profiles — Part 1: Preparation of chromatographic profiles for presentation in standards

ISO 11024-2, Essential oils — General guidance on chromatographic profiles — Part 2: Utilization of chromatographic profiles of samples of essential oils

#### 3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

#### 3.1

#### oil of rose

essential oil obtained by steam distillation of the flowers of  $Rosa \times damascena$  Miller, of the Rosaceae family, cultivated in Turkey, Morocco and Bulgaria

NOTE For information on the CAS number, see ISO/TR 21092.

#### 4 Requirements (all origins)

#### 4.1 Appearance

Liquid or more or less crystallized.

#### 4.2 Colour

Light yellow.

#### 4.3 Odour

Floral, rose.

#### 4.4 Relative density at 20 °C, $d_{20}^{20}$

Minimum: 0,848 Maximum: 0,880

#### 4.5 Refractive index at 20 °C

Minimum: 1,452 0 Maximum: 1,470 0

#### 4.6 Optical rotation at 20 °C

Between -5° and -1,8°.

#### Freezing point

Between +16 °C and +23,5 °C.

#### 4.8 Ester value

7 Minimum: Maximum: 24

#### 4.9 Chromatographic profile

Analysis of the essential oil shall be carried out by gas chromatography. In the chromatogram obtained, the representative and characteristic components shown in Table 1 shall be identified. The proportions of these components, indicated by the integrator, shall be as shown in Table 1. This constitutes the chromatographic profile of the essential oil.

#### 4.10 Flashpoint

Information on the flashpoint is given in Annex B.

#### Sampling

See ISO 212.

Minimum volume of test sample: 25 ml

This volume allows each of the tests specified in this International Standard to be carried out at least once.

#### Test methods

#### Relative density at 20 °C, $d_{20}^{20}$

See ISO 279.

#### 6.2 Refractive index at 20 °C

See ISO 280.

#### 6.3 Optical rotation at 20 °C

See ISO 592.

#### 6.4 Freezing point

See ISO 1041.

#### 6.5 Ester value

See ISO 709.

Time of saponification: 1 h

#### 6.6 Chromatographic profile

See ISO 11024-1 and ISO 11024-2.

#### Packaging, labelling, marking and storage

See ISO/TR 210 and ISO/TR 211.

Table 1 — Chromatographic profile

Values in percent

Component	Bulgaria		Turkey		Morocco		Turkey ("peasant" type)	
	min.	max.	min.	max.	min.	max.	min.	max.
Ethanol	_	2,0	_	7	_	3	_	2,0
Citronellol	20,0	34,0	34,0	49,0	30,0	47,0	26	40,0
Nerol	5,0	12,0	3,0	11,0	3,0	11,0	6,0	12,0
Geraniol	15,0	22,0	8,0	20,0	6,0	23,0	12,0	29,0
β-Phenylethanol	_	3,5	_	3,0	_	3,0	_	3,0
Heptadecane (Paraffin C <sub>17</sub> )	1,0	2,5	0,8	3,0	0,6	4,0	0,7	3,0
Nonadecane (Paraffin C <sub>19</sub> )	8,0	15,0	6,0	13,0	7,0	16,0	6,0	8,5
Heneicosane (Paraffin C <sub>21</sub> )	3,0	5,5	2,0	4,0	2,0	5,5	1,5	4,0

# Annex A

(informative)

# Typical chromatograms of the analysis by gas chromatography of the essential oil of rose ( $Rosa \times damascena$ Miller)

NOTE Only chromatograms taken on polar column are proposed, as on an apolar column nerol is coeluted with citronellol.

# Operating conditions

Column: silica capillary; length 60 m; internal diameter 0,32 mm

Stationary phase: bonded poly(ethylene glycol) (INNOWAX $^{\otimes}$ )

Heptadecane (Paraffin C<sub>17</sub>)

Citronellol

Peak identification

Ethanol

Film thickness: 0,5 µm

Oven temperature: isothermal at 80 °C for 5 min, then temperature programming from 80 °C to 220 °C at a rate of 2 °C/min

Injector temperature: 250 °C

Detector temperature: 250 °C

(Z)-9-Nonadecene (Paraffin C'<sub>19</sub>)

Nonadecane (Paraffin C<sub>19</sub>)

Geraniol

9

Nerol

4

Detector: flame ionization type Carrier gas: nitrogen

Volume injected: 0,2 µl

Carrier gas flow rate: 0,5 ml/min Split ratio: 1/30

Heneicosane (Paraffin C<sub>21</sub>)

eta-Phenylethanol Eicosane (Paraffin  $C_{20})$ 

Methyleugenol

Figure A.1 — Typical chromatogram taken on a polar column (origin: Turkey)

#### **Peak identification**

1 Ethanol

2 cis-Rose oxide

3 trans-Rose oxide

4 Linalol

5 Heptadecane (Paraffin C<sub>17</sub>)

6 Citronellol7 Nerol8 Geraniol

9 β-Phenylethanol

10 Nonadecane (Paraffin  $C_{19}$ )

11 n-Nonadecene (Alkene n-C $'_{19}$ )

12 Methyleugenol

13 Eicosane (Paraffin C<sub>20</sub>)

14 Alkene *n*-C<sub>20</sub>

15 Heneicosane (Paraffin C<sub>21</sub>)

16 Alkene *n*-C<sub>21</sub>
17 Eugenol
18 Alkane *n*-23

19 (E,E)-Farnesol

#### **Operating conditions**

Column: capillary; length 50 m; internal diameter 0,2 mm Stationary phase: poly(ethylene glycol) (CARBOWAX 20 M®)

Film thickness: 0,25 µm

Oven temperature: temperature programming from 65 °C to 230 °C at a rate of 2 °C/min

Injector temperature: 230 °C Detector temperature: 230 °C Detector flame ionization type

Carrier gas: hydrogen Volume injected: 0,2 µl

Carrier gas flow rate: 1,1 ml/min

Split ratio: 100/1

Figure A.2 — Typical chromatogram taken on a polar column (origin: Bulgaria)

## Annex B

(informative)

#### **Flashpoint**

#### **B.1 General information**

For safety reasons, transport companies, insurance companies, and people in charge of safety services require information on the flashpoints of essential oils, which in most cases are flammable products.

A comparative study on the relevant methods of analysis (see ISO/TR 11018) concluded that it was difficult to recommend a single apparatus for standardization purposes, given that:

- there is wide variation in the chemical composition of essential oils;
- the volume of the sample needed for certain requirements would be too costly for highpriced essential oils;
- as there are several different types of equipment which can be used for the determination, users cannot be expected to use one specified type only.

Consequently, it was decided to give a mean value for the flashpoint in an informative annex to each International Standard, for information, in order to meet the requirements of the interested parties.

The equipment with which this value was obtained should be specified.

For further information see ISO/TR 11018.

#### B.2 Flashpoint of oil of rose

The flashpoint should be confirmed with every batch of rose oil, as the content of ethanol is responsible for the flashpoint of this oil. The following values are given for information and they can vary between +37 °C (for an ethanol content of about 7 %) to +67 °C (for an ethanol content of about 2,5 %).

The mean value for the "peasant" essential oil is +37 °C (for an ethanol content of maximum 2 %).

NOTE Obtained with "Pensky Martens" equipment.

## **Bibliography**

- [1] ISO/TR 21092<sup>1)</sup>, Essential oils Characterization
- [2] ISO/TR 11018:1997, Essential oils General guidance on the determination of flashpoint

<sup>1)</sup> To be published.

ISO 9842:2003(E)

ICS 71.100.60

Price based on 7 pages